Dynamics of Cytoskeletons

Cell Biology and Molecular Biology
Transgenic and Mutant Mice for Neurological
diseases

B692515
Chung-Liang Chien (錢宗良)
Department of Anatomy and Cell Biology
College of Medicine, National Taiwan University
Taipei, Taiwan

Cytoskeletons:

Actin filaments Microtubules Intermediate filaments

- 1. Microtubule associated proteins: MAP2, Tau, and MAP1A
- 2. intermediate filaments: Neurofilament triplet proteins (NF-L, NF-M, and NF-H) peripherin, and a-internexin


ACTIN FILAMENTS



100 nm











Actin filaments (also known as microfilaments) are two-stranded helical polymers of the protein actin. They appear as flexible structures, with a diameter of 5-9 nm, and they are organized into a variety of linear bundles, two-dimensional networks, and three-dimensional gels. Although actin filaments are dispersed throughout the cell, they are most highly concentrated in the cortex, just beneath the plasma membrane.

Micrographs courtesy of Roger Craig (i and iv); P.T. Matsudaira and D.R. Burgess (ii); Keith Burridge (iii).


MICROTUBULES















Micrographs courtesy of Richard Wade (i); D.T. Woodrow and R.W. Linck (ii); David Shima (iii); A. Desai (iv).

Microtubules are long, hollow cylinders made of the protein tubulin. With an outer diameter of 25 nm, they are much more rigid than actin filaments.

Microtubules are long and straight and typically have one end attached to a single microtubule-organizing center (MTOC) called a centrosome, as shown


INTERMEDIATE FILAMENTS




100 nm













intermediate filaments are ropelike fibers with a diameter of around 10 nm; they are made of intermediate filament proteins, which constitute a large and heterogeneous family. One type of intermediate filament forms a meshwork called the nuclear lamina just beneath the inner nuclear membrane. Other types extend across the cytoplasm, giving cells mechanical strength. In an epithelial tissue, they span the cytoplasm from one cell-cell junction to another, thereby strengthening the entire epithelium.

Micrographs courtesy of Roy Quinlan (i); Nancy L. Kedersha (ii); Mary Osborn (iii); Ueli Aebi (iv).


Tagged α -internexin and EB-1 DNA constructs



(a gift from Professor Hirokawa, University of Tokyo)




Immunocytochemical staining of MAP1A transfected COS7 cells




DsRed-MAP1A + YFP-Tubulin in COS7 cell (Nocodazol treated)

10 second / frame, total 216 frames, 36 min




EB-1-YFP in COS7 cell 2 second / frame, total 151 frames, 5 min




EB-1-YFP in Neuro2A cell


10 second / frame, total 40 frames



Neural Differentiation of Mouse Embryonic Stem Cells




Neuronal differentiation from Embryoid Body in DMEM/F12 media supplemented with N2 for 5 days



Green: internexin Red: Vimentin

Intermediate Filament Proteins are good markers for determining the differentiation status of neural stem cells



Nestin, Vimentin, and Desmin

Internexin, Peripherin Neurofilament triplet Proteins (NF-L, NF-M, and NF-H)

Collaborators:

Nobutaka Hirokawa, University of Tokyo, Japan Norio Nakatsuji, Kyoto University, Japan Ronald Liem, Columbia University, U.S.A.

Lab. Members

Jin-Chung Shih (施景中) Pei Wang (王霈) Kwang-Wen Tseng (曾廣文) Tzu-Chiang Liu (劉自強)

Yi-Shan Lin (林憶珊) Yu-Chieh Lin (林雨潔) Ping-Chung Chen (陳品中)

