

Chapter 4: Higher-Order Differential Equations – Part 1

王奕翔

Department of Electrical Engineering
National Taiwan University

`ihwang@ntu.edu.tw`

October 10, 2013

Higher-Order Differential Equations

Most of this chapter deals with **linear** higher-order DE (except 4.10)

In our lecture, we skip 4.10 and focus on n -th order linear differential equations, where $n \geq 2$.

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

Methods of Solving Linear Differential Equations

We shall gradually fill up this slide as the lecture proceeds.

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

Initial-Value Problem (IVP)

An n -th order initial-value problem associate with (1) takes the form:

Solve:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

subject to:

$$y(x_0) = y_0, y'(x_0) = y_1, \dots, y^{(n-1)}(x_0) = y_{n-1} \quad (2)$$

Here (2) is a set of **initial conditions**.

Boundary-Value Problem (BVP)

Recall: in Chapter 1, we made 3 remarks on initial/boundary conditions

Remark (Initial vs. Boundary Conditions)

Initial Conditions: all conditions are at the **same** $x = x_0$.

Boundary Conditions: conditions can be at **different** x .

Remark (Number of Initial/Boundary Conditions)

“Usually” a n -th order ODE requires n initial/boundary conditions to specify an unique solution.

Remark (Order of the derivatives in the conditions)

Initial/boundary conditions can be the value or the function of 0-th to $(n - 1)$ -th order derivatives, where n is the order of the ODE.

Boundary-Value Problem (BVP)

Example (Second-Order ODE)

Consider the following second-order ODE

$$a_2(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_0(x) y = g(x) \quad (3)$$

- IVP: solve (3) s.t. $y(x_0) = y_0$, $y'(x_0) = y_1$.
- BVP: solve (3) s.t. $y(a) = y_0$, $y(b) = y_1$.
- BVP: solve (3) s.t. $y'(a) = y_0$, $y(b) = y_1$.
- BVP: solve (3) s.t.
$$\begin{cases} \alpha_1 y(a) + \beta_1 y'(a) = \gamma_1 \\ \alpha_2 y(b) + \beta_2 y'(b) = \gamma_2 \end{cases}$$

Existence and Uniqueness of the Solution to an IVP

Solve

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

subject to

$$y(x_0) = y_0, \quad y'(x_0) = y_1, \dots, y^{(n-1)}(x_0) = y_{n-1} \quad (2)$$

Theorem

If $a_n(x), a_{n-1}(x), \dots, a_0(x)$ and $g(x)$ are all continuous on an interval I , $a_n(x) \neq 0$ is not a zero function on I , and the initial point $x_0 \in I$, then the above IVP has a unique solution in I .

Existence and Uniqueness of the Solution to an IVP

Solve

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

subject to

$$y(x_0) = y_0, \quad y'(x_0) = y_1, \dots, \quad y^{(n-1)}(x_0) = y_{n-1} \quad (2)$$

Throughout this lecture, we assume that on some common interval I ,

- $a_n(x), a_{n-1}(x), \dots, a_0(x)$ and $g(x)$ are all continuous
- $a_n(x)$ is not a zero function, that is, $\exists x \in I$ such that $a_n(x) \neq 0$.

Existence and Uniqueness of the Solution to an BVP

Note: Unlike an IVP, even the n -th order ODE (1) satisfies the conditions in the previous theorem, a BVP corresponding to (1) may have many, one, or no solutions.

Example

Consider the 2nd-order ODE $\frac{d^2 y}{dx^2} + y = 0$, whose general solution takes the form $y = c_1 \cos x + c_2 \sin x$. Find the solution(s) to an BVP subject to the following boundary conditions respectively

- $y(0) = 0, y(2\pi) = 0$ Plug it in $\implies c_1 = 0, c_1 = 0$
 $\implies c_2$ is arbitrary \implies infinitely many solutions!
- $y(0) = 0, y(\pi/2) = 0$ Plug it in $\implies c_1 = 0, c_2 = 0$
 $\implies c_1 = c_2 = 0 \implies$ a unique solution!
- $y(0) = 0, y(2\pi) = 1$ Plug it in $\implies c_1 = 0, c_1 = 1$
 \implies contradiction \implies no solutions!

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

Homogeneous Equation

Linear n -th order ODE takes the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

Homogeneous Equation: $g(x)$ in (1) is a zero function:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0 \quad (4)$$

Nonhomogeneous Equation: $g(x)$ in (1) is **not** a zero function. Its *associated homogeneous equation* (4) is the one with the same coefficients except that $g(x)$ is a zero function

Later in the lecture we will see, when solving a nonhomogeneous equation, we must first solve its associated homogeneous equation (4).

Differential Operators

We introduce a **differential operator** D , which simply represent the operation of taking an ordinary differentiation:

Differential Operator

For a function $y = f(x)$, the differential operator D transforms the function $f(x)$ to its first-order derivative: $Dy := \frac{dy}{dx}$.

Higher-order derivatives can be represented compactly with D as well:

$$\frac{d^2 y}{dx^2} = D(Dy) =: D^2 y, \quad \frac{d^n y}{dx^n} =: D^n y$$

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y =: \left\{ \sum_{i=0}^n a_i(x) D^i \right\} y$$

Differential Operators and Linear Differential Equations

Note: Polynomials of differential operators are differential operators.

Let $L := \sum_{i=0}^n a_i(x)D^i$ be an n -th order differential operator.

Then we can compactly represent the linear differential equation (1) and the homogeneous linear DE (4) as

$$L(y) = g(x), \quad L(y) = 0$$

respectively.

Linearity and Superposition Principle

$L := \sum_{i=0}^n a_i(x)D^i$ is a **linear operator**: for two functions $f_1(x), f_2(x)$,

$$L(\lambda_1 f_1 + \lambda_2 f_2) = \lambda_1 L(f_1) + \lambda_2 L(f_2).$$

For any homogeneous linear equation (4), that is, $L(y) = 0$, we obtain the following superposition principle.

Theorem (Superposition Principle: Homogeneous Equations)

Let f_1, f_2, \dots, f_k be solutions to the homogeneous n -th order linear equation $L(y) = 0$ on an interval I , that is,

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0, \quad (4)$$

then the linear combination $f = \sum_{i=1}^k \lambda_i f_i$ is also a solution to (4).

Linear Dependence and Independence of Functions

In Linear Algebra, we learned that one can view the collection of all *functions* defined on a common interval as a **vector space**, where linear dependence and independence can be defined respectively.

Definition (Linear Dependence and Independence)

A set of functions $\{f_1(x), f_2(x), \dots, f_n(x)\}$ are **linearly dependent** on an interval I if $\exists c_1, c_2, \dots, c_n$ not all zero such that

$$c_1 f_1(x) + c_2 f_2(x) + \dots + c_n f_n(x) = 0, \quad \forall x \in I,$$

that is, the linear combination is a zero function. If the set of functions is not linearly dependent, it is **linearly independent**.

Example:

- $f_1(x) = \sin^2 x$, $f_2(x) = \cos^2 x$, $I = (-\pi, \pi)$: Linearly dependent
- $f_1(x) = 1$, $f_2(x) = x$, $f_3(x) = x^3$, $I = \mathbb{R}$: Linearly independent.

Linear Independence of Solutions to (4)

Consider the homogeneous linear n -th order DE

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0, \quad (4)$$

Given n solutions $\{f_1(x), f_2(x), \dots, f_n(x)\}$, we would like to test if they are independent or not.

Of course we can always go back to the definition but it is clumsy...

Recall: In Linear Algebra, to test if n vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ are linearly independent, we can compute the determinant of the matrix

$$\mathbf{V} := [\mathbf{v}_1 \quad \mathbf{v}_2 \quad \cdots \quad \mathbf{v}_n].$$

If $\det \mathbf{V} = 0$, they are linearly dependent; if $\det \mathbf{V} \neq 0$, they are linearly independent.

Criterion of Linearly Independent Solutions

Consider the homogeneous linear n -th order DE

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x) y = 0, \quad (4)$$

To test the linear independence of n solutions $\{f_1(x), f_2(x), \dots, f_n(x)\}$ to (4), we can use the following theorem.

Theorem

Let $\{f_1(x), f_2(x), \dots, f_n(x)\}$ be n solutions to the homogeneous linear n -th order DE (4) on an interval I . They are **linearly independent** on I

$$\iff W(f_1, f_2, \dots, f_n) := \begin{vmatrix} f_1 & f_2 & \cdots & f_n \\ f_1' & f_2' & \cdots & f_n' \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(n-1)} & f_2^{(n-1)} & \cdots & f_n^{(n-1)} \end{vmatrix} \neq 0.$$

Fundamental Set of Solutions

We are interested in describing the *solution space*, that is, the subspace spanned by the solutions to the homogeneous linear n -th order DE

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0. \quad (4)$$

How?

Recall: In Linear Algebra, we describe a subspace by its *basis*: any vector in the subspace can be represented by a linear combination of the elements in the basis, and these elements are linearly independent.

Similar things can be done here.

Definition (Fundamental Set of Solutions)

Any set $\{f_1(x), f_2(x), \dots, f_n(x)\}$ of n linearly independent solutions to the homogeneous linear n -th order DE (4) on an interval I is called a **fundamental set of solutions**.

General Solutions to Homogeneous Linear DE

General solution to an n -th order ODE:

An n -parameter family of solutions that can contains *all* solutions.

Theorem

Let $\{f_1(x), f_2(x), \dots, f_n(x)\}$ be a fundamental set of solutions to the homogeneous linear n -th order DE (4) on an interval I . Then the **general solution** to (4) is

$$y = c_1 f_1(x) + c_2 f_2(x) + \cdots + c_n f_n(x),$$

where $\{c_i \mid i = 1, 2, \dots, n\}$ are arbitrary constants.

Examples

Example

Consider the DE

$$\frac{d^2 y}{dx^2} = y.$$

Check that both $y = e^x$ and $y = e^{-x}$ are solutions to the equation. Derive the general solution to the DE.

A: The linear DE is homogeneous.

We see that $\frac{d^2}{dx^2} e^x = \frac{d}{dx} e^x = e^x$, and $\frac{d^2}{dx^2} e^{-x} = \frac{d}{dx} -e^{-x} = e^{-x}$. Hence they are both solutions to the homogeneous linear second-order DE.

Since

$$\begin{vmatrix} e^x & e^{-x} \\ e^x & -e^{-x} \end{vmatrix} = -1 - 1 = -2 \neq 0,$$

the two solutions are linearly independent. Hence, the general solution can be written as $y = c_1 e^x + c_2 e^{-x}$, $c_1, c_2 \in \mathbb{R}$.

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - **Nonhomogeneous Equations**
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

General Solutions to Nonhomogeneous Linear DE

Nonhomogeneous linear n -th order ODE takes the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

or equivalently, $L(y) = g(x)$, $L := \sum_{i=0}^n a_i(x) D^i$

where $g(x)$ is not a zero function.

How to find its general solution?

Idea:

- Find the general solution y_c to the *homogeneous* equation $L(y) = 0$.
- Find **a** solution y_p to the *nonhomogeneous* equation $L(y) = g(x)$.
- The general solution $y = y_c + y_p$.

General Solutions to Nonhomogeneous Linear DE

Nonhomogeneous :

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x) \quad (1)$$

Homogeneous :

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0 \quad (4)$$

Theorem

Let y_p be any particular solution to the nonhomogeneous linear n -th order DE (1) on an interval I , and y_c be the general solution to the associated homogeneous linear n -th order DE (4) on I , then the general solution to (1) is

$$y = y_c + y_p.$$

Proof of the Theorem

Proof: Let $y = f(x)$ be any solution to the nonhomogeneous linear n -th order DE (1), that is, $L(y) = g(x)$.

Now, since both y_p and f are solutions to $L(y) = g(x)$, we have

$$0 = L(f) - L(y_p) = L(f - y_p).$$

Hence, $(f - y_p)$ is a solution to the homogeneous linear n -th order DE (4).

Therefore, any solution to (1) can be represented by the sum of a solution to (4) and the particular solution y_p .

Examples

Example

Consider the DE

$$\frac{d^2 y}{dx^2} = y + 9.$$

Derive the general solution to the DE.

A: The linear DE is nonhomogeneous. The associated homogeneous equation $\frac{d^2 y}{dx^2} = y$ has the following general solution:

$$y = c_1 e^x + c_2 e^{-x}, \quad c_1, c_2 \in \mathbb{R}.$$

There is an obvious particular solution $y = -9$.

Hence, the general solution can be written as

$$y = c_1 e^x + c_2 e^{-x} - 9, \quad c_1, c_2 \in \mathbb{R}$$

Superposition Principle for Nonhomogeneous Equations

For nonhomogeneous linear differential equations, we have the following superposition principle.

Theorem (Superposition Principle: Nonhomogeneous Equations)

Let $f_i(x)$ be a particular solution to the nonhomogeneous n -th order linear equation $L(y) = g_i(x)$ on an interval I , for $i = 1, 2, \dots, k$. Then the linear combination $f = \sum_{i=1}^k \lambda_i f_i$ is a particular solution to the nonhomogeneous n -th order linear equation

$$L(y) = \sum_{i=1}^k \lambda_i g_i(x).$$

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

Finding a New Solution

Recall: the fundamental set of solutions of the homogeneous linear n -th order DE

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0 \quad (4)$$

contains n linearly independent solutions.

Now suppose we already have k ($1 \leq k < n$) linearly independent solutions $\{f_1, f_2, \dots, f_k\}$. How do we find another one f_{k+1} so that the $(k+1)$ solutions $\{f_1, f_2, \dots, f_{k+1}\}$ remain linearly independent?

Second Order Equation

We begin with the simplest case: $n = 2$ and $k = 1$. Consider the following homogeneous linear second order DE

$$a_2(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_0(x) y = 0.$$

Suppose we already have a solution $y = f_1(x)$. How do we find another solution $y = f_2(x)$, such that f_1 and f_2 are linearly independent?

Idea: Let $f_2(x) = u(x)f_1(x)$, and make use of the fact that

$$a_2(x) \frac{d^2}{dx^2} f_1 + a_1(x) \frac{d}{dx} f_1 + a_0(x) f_1 = 0$$

to reduce the second order DE into a **first order DE of u** !

Example

Example

$f_1(x) = x^2$ is a solution of the second order DE $x^2 \frac{d^2 y}{dx^2} - 3x \frac{dy}{dx} + 4y = 0$.
Find the general solution of the above DE for $x > 0$.

A: We need to find a fundamental set of solutions, which contains two linearly independent solutions. Now we have only one. To find a second one, let us set substitute $y = f_1 u = x^2 u$:

$$\frac{dy}{dx} = 2xu + x^2 u', \quad \frac{d^2 y}{dx^2} = (2u + 2xu') + (2xu' + x^2 u'') = 2u + 4xu' + x^2 u''$$

$$\begin{aligned} \implies x^2 \frac{d^2 y}{dx^2} - 3x \frac{dy}{dx} + 4y &= 2x^2 u + 4x^3 u' + x^4 u'' - 6x^2 u - 3x^3 u' + 4x^2 u \\ &= x^3 u' + x^4 u'' = 0 \end{aligned}$$

$$\implies v + xv' = 0 \quad (\text{Set } v := u')$$

$$\implies \text{one such } v = \frac{1}{x} \implies \text{one such } u = \ln x.$$

Example

Example

$f_1(x) = x^2$ is a solution of the second order DE $x^2 \frac{d^2 y}{dx^2} - 3x \frac{dy}{dx} + 4y = 0$.
Find the general solution of the above DE for $x > 0$.

We find a second solution $y = f_2(x) = x^2 \ln x$ on $x \in (0, \infty)$, and the general solution is

$$y = c_1 x^2 + c_2 x^2 \ln x.$$

Question:

How about the more complicated case, when $n > 2$ and $k > 1$?

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

In this section we focus on solving (that is, giving general solutions to)
Homogeneous Linear Equations with Constant Coefficients

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 \frac{dy}{dx} + a_0 y = 0, \quad (5)$$

which is a homogeneous linear DE with **constant real coefficients**.

In the textbook, it tells us (without much reasoning) what the form of the general solution should look like, and then we analyze the particular structure of a give equation to derive the exact form.

In this lecture, we try to provide more reasoning, so that you get a clearer big picture.

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

Second Order Equation

We begin with some examples of second order equations.

Example

Find the general solution of $\frac{d^2 y}{dx^2} - 3\frac{dy}{dx} + 2y = 0$.

A: Let us use the differential operator to rewrite this DE as follows:

$$(D^2 - 3D + 2)y = 0.$$

Note that $L := D^2 - 3D + 2 = (D - 1)(D - 2)$.

We can view the second-order differential operator L as a concatenation of two first-order differential operators: $(D - 1)$ and $(D - 2)$!

$$y \longrightarrow \boxed{L := D^2 - 3D + 2} \longrightarrow \frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y$$

|||

$$L = (D - 2)(D - 1)$$

$$y \longrightarrow \boxed{D - 1} \longrightarrow \boxed{D - 2} \longrightarrow \frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y$$

|||

$$L = (D - 1)(D - 2)$$

$$y \longrightarrow \boxed{D - 2} \longrightarrow \boxed{D - 1} \longrightarrow \frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y$$

Second Order Equation

Example

Find the general solution of $\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + 2y = 0$.

A: We have found the equivalent forms of the above equation

$$Ly = 0 \quad \equiv \quad (D - 2) \{(D - 1)y\} = 0 \quad \equiv \quad (D - 1) \{(D - 2)y\} = 0$$

where $L := D^2 - 3D + 2 = (D - 1)(D - 2)$.

Observation:

- If f_1 is a solution to $(D - 1)y = 0$, it is also a solution to $Ly = 0$.
A solution: $f_1 = e^x$.
- If f_2 is a solution to $(D - 2)y = 0$, it is also a solution to $Ly = 0$.
A solution: $f_2 = e^{2x}$.

Second Order Equation

Example

Find the general solution of $\frac{d^2 y}{dx^2} - 3\frac{dy}{dx} + 2y = 0$.

A: So far we have found two solutions to $(D^2 - 3D + 2)y = 0$:

$$f_1 = e^x, \quad \text{corresponds to } (D - 1)y = 0$$

$$f_2 = e^{2x}, \quad \text{corresponds to } (D - 2)y = 0.$$

f_1 and f_2 are linearly independent (**Exercise:** check!) and hence $\{f_1, f_2\}$ is a fundamental set of solutions.

\implies The general solution:

$$y = c_1 f_1 + c_2 f_2 = \boxed{c_1 e^x + c_2 e^{2x}}, \quad c_1, c_2 \in \mathbb{R}.$$

How we solve $\frac{d^2 y}{dx^2} - 3\frac{dy}{dx} + 2y = 0$

- 1 Use a polynomial of D ,

$$L := p(D) = D^2 - 3D + 2,$$

to rewrite the DE into the form $Ly = 0$.

- 2 Factor $p(D) = (D - 1)(D - 2)$.
- 3 Observe that a solution to either $(D - 1)y = 0$ or $(D - 2)y = 0$ will be a solution to $Ly = 0$.
- 4 Find two solutions $f_1 = e^x$ and $f_2 = e^{2x}$, corresponding to $(D - 1)y = 0$ and $(D - 2)y = 0$ respectively.
- 5 Check that f_1 and f_2 are linearly independent, and hence they form a fundamental set of solutions.
- 6 Finally we get the general solution $y = c_1 e^x + c_2 e^{2x}$.

$p(D) = a_2D^2 + a_1D + a_0$ Has Two Distinct Real Roots

For a homogeneous linear second order DE with constant coefficients
 $Ly = 0$, where (WLOG we assume $a_2 = 1$)

$$L := p(D) = a_2D^2 + a_1D + a_0 = D^2 + a_1D + a_0 :$$

Fact

If $p(D)$ has two distinct real roots m_1 and m_2 , then we can use the above mentioned method to get a general solution

$$y = c_1 e^{m_1 x} + c_2 e^{m_2 x}.$$

What if $p(D)$ has

- Two repeated real roots, or
- Two conjugate complex roots?

$p(D)$ Has Two Conjugate Complex Roots $\alpha \pm i\beta$

Suppose $p(D)$ has two conjugate complex roots

$$m_1 = \alpha + i\beta, \quad m_2 = \alpha - i\beta, \quad \alpha, \beta \in \mathbb{R}.$$

If we slightly extend our discussion to complex-valued DE, it is not hard to see that the previous method works again and we get a general (complex-valued) solution

$$y = C_1 e^{m_1 x} + C_2 e^{m_2 x}, \quad C_1, C_2 \in \mathbb{C}.$$

Still we need to get back to the real domain ...

So, let's do some further manipulation by using the fact that

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

$p(D)$ Has Two Conjugate Complex Roots $\alpha \pm i\beta$

The general solution to $Ly = 0$ where $L = p(D)$ is

$$\begin{aligned}y &= C_1 e^{(\alpha+i\beta)x} + C_2 e^{(\alpha-i\beta)x} = C_1 e^{\alpha x} e^{i\beta x} + C_2 e^{\alpha x} e^{-i\beta x} \\&= C_1 e^{\alpha x} (\cos \beta x + i \sin \beta x) + C_2 e^{\alpha x} (\cos \beta x - i \sin \beta x) \\&= (C_1 + C_2) e^{\alpha x} \cos \beta x + i(C_1 - C_2) e^{\alpha x} \sin \beta x\end{aligned}$$

To get a real-valued solution, there are two choices:

- Pick $C_1 + C_2 = 1$, $C_1 - C_2 = 0$: we get $y = f_1(x) = e^{\alpha x} \cos \beta x$.
- Pick $C_1 + C_2 = 0$, $C_1 - C_2 = -i$: we get $y = f_2(x) = e^{\alpha x} \sin \beta x$.

Since f_1 and f_2 are linearly independent, the general real-valued solution to $Ly = 0$ where $L = p(D)$ is

$$y = c_1 e^{\alpha x} \cos \beta x + c_2 e^{\alpha x} \sin \beta x, \quad c_1, c_2 \in \mathbb{R}.$$

$p(D)$ Has Two Repeated Real Roots m

Suppose $p(D)$ has two repeat real roots m , which means that

$$p(D) = (D - m)^2.$$

$$L = (D - m)^2$$

$$y \rightarrow \boxed{D - m} \rightarrow \boxed{D - m} \rightarrow \frac{d^2y}{dx^2} - 2m \frac{dy}{dx} + m^2y$$

From the previous discussion, we see that $y = f_1(x) = e^{mx}$ is a solution to $(D - m)y = 0$ and hence it is also a solution to $(D - m)^2y = 0$.

Question:

How to find another solution $y = f_2(x)$ so that f_1 and f_2 are linearly independent?

$p(D)$ Has Two Repeated Real Roots m

$f_1(x) = e^{mx}$ is a solution to $(D - m)^2 y = 0$ because:

$$f_1(x) = e^{mx} \longrightarrow \boxed{D - m} \xrightarrow{0} \boxed{D - m} \longrightarrow 0$$

Why not find some $f_2(x)$ such that after the first $\boxed{D - m}$ block, the outcome is $f_1(x) = e^{mx}$?

$$f_2(x) = ? \longrightarrow \boxed{D - m} \xrightarrow{e^{mx}} \boxed{D - m} \longrightarrow 0$$

We only need to solve a first order linear DE!

$$\underbrace{e^{-mx}}_{\text{integrating factor}} f_2(x) = \int \overbrace{\underbrace{e^{-mx}}_{\text{integrating factor}} e^{mx}}^1 dx \implies f_2(x) = xe^{mx}.$$

$p(D)$ Has Two Repeated Real Roots m

We have found two solutions to $(D - m)^2 y = 0$:

$$f_1(x) = e^{mx}, \quad f_2(x) = xe^{mx},$$

and they are linearly independent (check!).

Hence the general solution to $p(D)y = 0$ is

$$y = c_1 e^{mx} + c_2 x e^{mx} = \boxed{(c_1 + c_2 x) e^{mx}}.$$

Summary: Second Order Equation $a_2 \frac{d^2 y}{dx^2} + a_1 \frac{dy}{dx} + a_0 = 0$

Define the following (quadratic) polynomial

$$p(D) := a_2 D^2 + a_1 D + a_0.$$

Roots of $p(D)$	General Solution
Distinct real roots $m_1, m_2 \in \mathbb{R}$	$y = c_1 e^{m_1 x} + c_2 e^{m_2 x}$
Conjugate complex roots $\alpha \pm i\beta, \alpha, \beta \in \mathbb{R}$	$y = c_1 e^{\alpha x} \sin \beta x + c_2 e^{\alpha x} \cos \beta x$
Repeated real roots $m \in \mathbb{R}$	$y = (c_1 + c_2 x) e^{m x}$

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

n -th Order Equation $a_n \frac{d^n y}{dx^n} + \cdots + a_1 \frac{dy}{dx} + a_0 = 0$

Define

$$p(D) := a_n D^n + a_{n-1} D^{n-1} + \cdots + a_1 D + a_0 = \sum_{i=0}^n a_i D^i$$

and rewrite the n -th order equation as $\boxed{p(D)y = 0}$.

$p(D)$: a polynomial of order n with real-valued coefficients.

- $p(D)$ has n roots in the complex domain (counting the **multiplicity**)
- Complex roots of $p(D)$ must appear in conjugate pairs.

Example: $p(D) = (D - 1)^3(D - 2)^1(D^2 - 2D + 2)^2$ is a polynomial of order 8, and has the following roots

1	multiplicity 3
2	multiplicity 1
$1 \pm i$	multiplicity 2 for each.

Finding the General Solution of $p(D)y = 0$


High-level Idea: let $p(D)$ have n_1 distinct real roots $\{m_i \mid i \in [1 : n_1]\}$, and n_2 distinct pairs of conjugate complex roots $\{\alpha_j \pm i\beta_j \mid j \in [1 : n_2]\}$.

1 Factorize $p(D) = \sum_{i=0}^n a_i D^i$ as

$$p(D) = a_n \left(\prod_{i=1}^{n_1} \overbrace{(D - m_i)^{k_i}}^{p_i(D)} \right) \left(\prod_{j=1}^{n_2} \overbrace{(D^2 - 2\alpha_j D + \alpha_j^2 + \beta_j^2)^{l_j}}^{q_j(D)} \right)$$
$$= a_n \prod_{i=1}^{n_1} p_i(D) \prod_{j=1}^{n_2} q_j(D), \text{ where } n = \sum_{i=1}^{n_1} k_i + 2 \sum_{j=1}^{n_2} l_j.$$

- 2 For each $i \in [1 : n_1]$, find k_i linearly independent solutions of $p_i(D)y = 0$.
- 3 For each $j \in [1 : n_2]$, find $2l_j$ linearly independent solutions of $q_j(D)y = 0$.
- 4 Combine them all to get n linearly independent solutions of $p(D)y = 0$.

$$p(D) := \sum_{i=0}^n a_i D^i, \quad a_n = 1$$



$$p_i(D) := (D - m_i)^{k_i}, \quad i \in [1 : n_1]; \quad q_j(D) := (D^2 - 2\alpha_j D + \alpha_j^2 + \beta_j^2)^{l_j}, \quad j \in [1 : n_2].$$


$p(D)$ have n_1 distinct real roots $\{m_i \mid i \in [1 : n_1]\}$, and n_2 distinct pairs of conjugate complex roots $\{\alpha_j \pm i\beta_j \mid j \in [1 : n_2]\}$.

Note: The solutions of different blocks in the above diagram will be linearly independent.

Solve $(D - m)^k y = 0$

$k = 2$: two linearly independent solutions $f_1(x) = e^{mx}$ and $f_2(x) = xe^{mx}$.

$k = 3$: Look at the diagram below:



We only need to solve a first order linear DE!

$$\underbrace{e^{-mx}}_{\text{integrating factor}} f_3(x) = \int \overbrace{\underbrace{e^{-mx}}_{\text{integrating factor}} \underbrace{xe^{mx}}_x}_{\text{integrating factor}} dx \implies f_3(x) = x^2 e^{mx} / 2.$$

We can drop the factor of 2 and pick $f_3(x) = x^2 e^{mx}$.

Solve $(D - m)^k y = 0$

$$f_{i+1}(x) = ? \rightarrow \boxed{D - m} \xrightarrow{x^{i-1} e^{mx}} \boxed{(D - m)^i} \rightarrow 0$$

We can repeat this procedure and find k linearly independent solutions:

$$\boxed{f_1(x) = e^{mx}, f_2(x) = xe^{mx}, f_3(x) = x^2 e^{mx}, \dots, f_k(x) = x^{k-1} e^{mx}}.$$

$$\text{Solve } (D^2 - 2\alpha D + \alpha^2 + \beta^2)^l y = 0$$

$$D^2 - 2\alpha D + \alpha^2 + \beta^2 = (D - m)(D - \bar{m}), \text{ where } m = \alpha + i\beta \in \mathbb{C}.$$

$$\therefore (D^2 - 2\alpha D + \alpha^2 + \beta^2)^l = (D - m)^l (D - \bar{m})^l$$

We can repeat the previous discussion and get $2l$ linearly independent solutions (in \mathbb{C}):

$$F_1(x) = e^{mx}, F_2(x) = xe^{mx}, \dots, F_l(x) = x^{l-1} e^{mx}$$

$$\bar{F}_1(x) = e^{\bar{m}x}, \bar{F}_2(x) = xe^{\bar{m}x}, \dots, \bar{F}_l(x) = x^{l-1} e^{\bar{m}x}$$

For each $j \in [1 : l]$, use F_j and \bar{F}_j to generate two real-valued solutions:

$$f_{2j-1}(x) = \frac{1}{2}F_j(x) + \frac{1}{2}\bar{F}_j(x) = \text{Re}\{F_j(x)\} = x^{j-1} e^{\alpha x} \cos \beta x$$

$$f_{2j}(x) = \frac{-i}{2}F_j(x) + \frac{i}{2}\bar{F}_j(x) = \text{Im}\{F_j(x)\} = x^{j-1} e^{\alpha x} \sin \beta x.$$

$$\text{Solve } (D^2 - 2\alpha D + \alpha^2 + \beta^2)^l y = 0$$

Here are $2l$ linearly independent real-valued solutions:

$$\left\{ x^{j-1} e^{\alpha x} \cos \beta x, x^{j-1} e^{\alpha x} \sin \beta x \mid j = 1, 2, \dots, l \right\}$$

Examples

Example

Solve the IVP $4y'' + 4y' + 17y = 0$, $y(0) = -1$, $y'(0) = 2$.

A: Write down the associated polynomial

$$p(D) = 4D^2 + 4D + 17 = (2D + 1)^2 + 16.$$

$p(D)$ has two roots: $-\frac{1}{2} \pm 2i$. Hence according to the previous discussion, the general solution is $y = c_1 f_1(x) + c_2 f_2(x)$, where

$$f_1(x) = e^{-\frac{1}{2}x} \cos 2x, \quad f_2(x) = e^{-\frac{1}{2}x} \sin 2x.$$

Plug in the initial conditions, we get

$$\begin{cases} c_1 f_1(0) + c_2 f_2(0) = -1 \\ c_1 f_1'(0) + c_2 f_2'(0) = 2 \end{cases} \implies c_1 = -1, c_2 = \frac{3}{4}$$

Examples

Example

Solve the IVP $y''' + 2y'' - 2y = 0$, $y(0) = 1$, $y'(0) = 2$, $y''(0) = 4$.

A: Write down the associated polynomial

$$p(D) = D^3 + 2D^2 - 2 = (D - 1)(D^2 + 2D + 2).$$

$p(D)$ has three roots: $1, -1 \pm i$. Hence according to the previous discussion, the general solution is $y = c_1 f_1(x) + c_2 f_2(x) + c_3 f_3(x)$, where

$$f_1(x) = e^x, \quad f_2(x) = e^{-x} \cos x, \quad f_3(x) = e^{-x} \sin x.$$

Plug in the initial conditions, we get

$$\begin{cases} c_1 f_1(0) + c_2 f_2(0) + c_3 f_3(0) = 1 \\ c_1 f_1'(0) + c_2 f_2'(0) + c_3 f_3'(0) = 2 \\ c_1 f_1''(0) + c_2 f_2''(0) + c_3 f_3''(0) = 4 \end{cases} \implies c_1 = 2, c_2 = -1, c_3 = -1.$$

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

Homogeneous Linear DE with Variable Coefficients

In general, a homogeneous linear differential equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0 \quad (4)$$

may have *variable* coefficients, that is, $a_i(x)$ is not a constant function, for $i = 0, 1, \dots, n$.

If the coefficients are not constants, it is usually hard to find closed-form solutions. Instead, we shall see in Chapter 6 that the best we expect is to find a solution in the form of *infinite series*.

There is one exception: when

$$a_i(x) = a_i x^i, \quad \forall i = 0, 1, \dots, n.$$

Cauchy-Euler Equation

Leonhard Euler



Augustin-Louis Cauchy



Cauchy-Euler Equation

Definition (Cauchy-Euler Equation)

A linear differential equation of the form

$$a_n x^n \frac{d^n y}{dx^n} + a_{n-1} x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 x \frac{dy}{dx} + a_0 y = g(x)$$

is called a Cauchy-Euler equation.

We first focus on finding the general solution of a **homogeneous** Cauchy-Euler equation (see below) in this lecture. In the next lecture we discuss how to find a particular solution of a nonhomogeneous Cauchy-Euler equation.

$$a_n x^n \frac{d^n y}{dx^n} + a_{n-1} x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 x \frac{dy}{dx} + a_0 y = 0$$

Change of Variable

Goal: Find the general solution of a homogeneous Cauchy-Euler equation on the interval $(0, \infty)$.

$$a_n x^n \frac{d^n y}{dx^n} + a_{n-1} x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 x \frac{dy}{dx} + a_0 y = 0$$

Idea: Convert a Cauchy-Euler equation into a linear equation with constant coefficients, by substituting $x = e^t \implies \frac{dx}{dt} = e^t = x$.

Observation

With the substitution $x = e^t \implies \frac{dx}{dt} = e^t = x$,

$$\frac{dy}{dt} = \frac{dx}{dt} \frac{dy}{dx} = x \frac{dy}{dx} \implies x \frac{dy}{dx} = \frac{dy}{dt} := D_t y$$

$$D_t^2 y = \frac{dx}{dt} \frac{d}{dx} \left(x \frac{dy}{dx} \right) = x \left(\frac{dy}{dx} + x \frac{d^2 y}{dx^2} \right) = D_t y + x^2 \frac{d^2 y}{dx^2}$$

$$\implies x^2 \frac{d^2 y}{dx^2} = D_t(D_t - 1)y$$

$$D_t^2(D_t - 1)y = \frac{dx}{dt} \frac{d}{dx} \left(x^2 \frac{d^2 y}{dx^2} \right) = x \left(2x \frac{d^2 y}{dx^2} + x^2 \frac{d^3 y}{dx^3} \right)$$

$$= 2D_t(D_t - 1)y + x^3 \frac{d^3 y}{dx^3}$$

$$\implies x^3 \frac{d^3 y}{dx^3} = D_t(D_t - 1)(D_t - 2)y$$

Conversion

Fact

With $x = e^t$, $x^k D_x^k = D_t(D_t - 1) \cdots (D_t - k + 1)$, for all integer $k \geq 1$.

Proof: For $k = 1$, the fact is true. Suppose the fact holds for $k = h \geq 1$. Then we have $x^h D_x^h y = D_t(D_t - 1) \cdots (D_t - h + 1) y$.

Take the derivative with respect to x on both sides, we get

$$\begin{aligned} (hx^{h-1} D_x^h + x^h D_x^{h+1}) y &= \frac{dt}{dx} D_t^2 (D_t - 1) \cdots (D_t - h + 1) y \\ \implies (hx^h D_x^h + x^{h+1} D_x^{h+1}) y &= D_t^2 (D_t - 1) \cdots (D_t - h + 1) y \\ \implies (hD_t(D_t - 1) \cdots (D_t - h + 1) + x^{h+1} D_x^{h+1}) y &= D_t^2 (D_t - 1) \cdots (D_t - h + 1) y \\ \implies x^{h+1} D_x^{h+1} &= D_t(D_t - 1) \cdots (D_t - h + 1)(D_t - h) \end{aligned}$$

Hence we prove the fact by induction.

Convert into a Equation with Constant Coefficients

Based on the above fact, with the substitution $x = e^t$, we convert a Cauchy-Euler Equation

$$a_n x^n \frac{d^n y}{dx^n} + a_{n-1} x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 x \frac{dy}{dx} + a_0 y = 0$$

into a linear differential equation (with respect to t) with constant coefficients

$$L_t y = 0, \quad L_t := \sum_{i=0}^n a_i D_t (D_t - 1) \cdots (D_t - i + 1)$$

Mapping of Solutions: with $x = e^t$,

t Domain	$t^k e^{mt}$	$t^k e^{\alpha t} \cos \beta t$	$t^k e^{\alpha t} \sin \beta t$
x Domain	$(\ln x)^k x^m$	$(\ln x)^k x^\alpha \cos(\beta \ln x)$	$(\ln x)^k x^\alpha \sin(\beta \ln x)$

Solutions for $x < 0$

So far we give the general solution of the Cauchy-Euler equation for $x \in (0, \infty)$. How about the solution for $x < 0$?

Idea: Change of variable – substitute $x = -u$, and solve the new Cauchy-Euler Equation for $u > 0$.

Conversion:

$$D_x^k = (-1)^k D_u^k.$$

- 1 Preliminary: Linear Equations
 - Initial-Value and Boundary-Value Problems
 - Homogeneous Equations
 - Nonhomogeneous Equations
- 2 Reduction of Order
- 3 Homogeneous Linear Equations with Constant Coefficients
 - Second Order Equations
 - n -th Order Equations
- 4 Cauchy-Euler Equation
- 5 Summary

Short Recap

- Initial-Value Problems (IVP) vs. Boundary-Value Problems (BVP)
- Homogeneous vs Nonhomogeneous Linear ODE
- Fundamental set of solutions and General Solutions
- Linearity and Superposition Principle
- General Solution of Homogeneous Linear Equation with Constant Coefficients: Usage of Polynomial of Differential Operator D
- General Solution of Homogeneous Cauchy-Euler Equation:
Substitution $x = e^t$ and $x^k D_x^k = \prod_{i=1}^k (D_t - i + 1)$

Self-Practice Exercises

4-1: 1, 9, 13, 17, 21, 25, 35

4-2: 1, 3, 13, 17, 19

4-3: 3, 5, 17, 21, 25, 31, 37, 51, 57

4-7: 1, 5, 15, 25, 31, 41