

Chapter 7: The Laplace Transform – Part 2

王奕翔

Department of Electrical Engineering
National Taiwan University

ihwang@ntu.edu.tw

November 26, 2013

So far we have learned

- 1 Basic properties of Laplace Transform
- 2 Inverse transform ([Memorize with Laplace transform in pairs!](#))
- 3 How to use Laplace Transform to solve an IVP

End of story?

Solving a Second-Order IVP with Laplace Transform

Example

Solve $y'' - 2y' + y = e^{2t}$, $y(0) = 1$, $y'(0) = 5$.**Step 1:** Laplace-transform both sides:

$$\begin{aligned} \mathcal{L}\{y''\} - 2\mathcal{L}\{y'\} + \mathcal{L}\{y\} &= \mathcal{L}\{e^{2t}\} \\ \implies (s^2 Y(s) - sy(0) - y'(0)) - 2(sY(s) - y(0)) + Y(s) &= \frac{1}{s-2} \\ \implies (s^2 - 2s + 1)Y(s) &= s + 3 + \frac{1}{s-2} \end{aligned}$$

Step 2: Solve $Y(s)$: $Y(s) = \frac{s+3}{(s-1)^2} + \frac{1}{(s-1)^2(s-2)}$.**Step 3:** Compute the inverse Laplace transform of $Y(s)$: [How to compute?](#)

$$Y(s) = \frac{3}{(s-1)^2} + \frac{1}{s-2} \implies y(t) = 3\mathcal{L}^{-1}\left\{\frac{1}{(s-1)^2}\right\} + e^{2t}.$$

Partial fraction decomposition:

$$\frac{s+3}{(s-1)^2} = \frac{(s-1)+4}{(s-1)^2} = \frac{1}{s-1} + \frac{4}{(s-1)^2}.$$

$$\frac{1}{(s-1)^2(s-2)} = \frac{A}{s-2} + \frac{B}{s-1} + \frac{C}{(s-1)^2}$$

$$A = \left[\frac{1}{(s-1)^2 \cancel{(s-2)}} \right]_{s=2} = 1$$

$$C = \left[\frac{1}{\cancel{(s-1)}^2 (s-2)} \right]_{s=1} = -1$$

$$1 = (B(s-1) + C)(s-2) + A(s^2 - 2s + 1) \implies B = -A = -1.$$

We already know that $\mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} = t$.

If we know what is the inverse transform of a function $F(s)$ when it is **translated** by 1 in the s -axis, that is, $\mathcal{L}^{-1} \{F(s - 1)\}$, we can solve!

Need more properties of Laplace and its inverse transforms!

1 Properties of Translations

Translation on the s -Axis

Recall that

$$1 \xrightarrow{\mathcal{L}} \frac{1}{s}, \quad e^{at} \xrightarrow{\mathcal{L}} \frac{1}{s-a}.$$

Multiplying 1 by e^{at} in t -domain results in **right-shift of a** in s -domain.


Theorem

Let $f(t) \xrightarrow{\mathcal{L}} F(s)$. For any a ,

$$\mathcal{L} \{ e^{at} f(t) \} = F(s - a).$$

Proof:

$$\mathcal{L} \{ e^{at} f(t) \} = \int_0^{\infty} e^{-st} e^{at} f(t) dt = \int_0^{\infty} e^{-(s-a)t} f(t) dt = F(s - a).$$



Back to the Problem

Example

Solve $y'' - 2y' + y = e^{2t}$, $y(0) = 1$, $y'(0) = 5$.

Step 3: Compute the inverse Laplace transform of $Y(s)$: [How to compute?](#)

$$Y(s) = \frac{3}{(s-1)^2} + \frac{1}{s-2} \implies y(t) = 3\mathcal{L}^{-1}\left\{\frac{1}{(s-1)^2}\right\} + e^{2t}.$$

$$\mathcal{L}^{-1}\left\{\frac{1}{(s-1)^2}\right\} = e^t \mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} = te^t.$$

Hence,

$$Y(s) = \frac{3}{(s-1)^2} + \frac{1}{s-2} \implies \boxed{y(t) = 3te^t + e^{2t}}.$$

Laplace Transform of $t^n e^{at}$

$$\mathcal{L} \{t^n e^{at}\} = \frac{n!}{(s-a)^{n+1}}, \quad n = 0, 1, 2, \dots, \quad s > a$$

We can obtain the inverse Laplace transform of $\frac{1}{(s-a)^n}$:

$$\mathcal{L}^{-1} \left\{ \frac{1}{(s-a)^n} \right\} = \frac{t^{n-1}}{(n-1)!} e^{at}, \quad n = 1, 2, \dots$$

Laplace Transform of $e^{at} \sin(kt)$ and $e^{at} \cos(kt)$

$$\mathcal{L} \{ e^{at} \sin(kt) \} = \frac{k}{(s-a)^2 + k^2}, \quad s > a$$

$$\mathcal{L} \{ e^{at} \cos(kt) \} = \frac{(s-a)}{(s-a)^2 + k^2}, \quad s > a$$

We can obtain the corresponding inverse Laplace transforms:

$$\mathcal{L}^{-1} \left\{ \frac{k}{(s-a)^2 + k^2} \right\} = e^{at} \sin(kt)$$

$$\mathcal{L}^{-1} \left\{ \frac{(s-a)}{(s-a)^2 + k^2} \right\} = e^{at} \cos(kt)$$

Solving a Second-Order IVP with Laplace Transform

Example

Solve $y'' - 4y' + 5y = t^2 e^{2t}$, $y(0) = 2$, $y'(0) = 6$.**Step 1:** Laplace-transform both sides:

$$\begin{aligned} \mathcal{L}\{y''\} - 4\mathcal{L}\{y'\} + 5\mathcal{L}\{y\} &= \mathcal{L}\{t^2 e^{2t}\} \\ \implies (s^2 Y(s) - sy(0) - y'(0)) - 4(sY(s) - y(0)) + 5Y(s) &= \frac{2}{(s-2)^3} \\ \implies (s^2 - 4s + 5)Y(s) &= 2s - 2 + \frac{2}{(s-2)^3} \end{aligned}$$

Step 2: Solve $Y(s)$: $Y(s) = \frac{2s-2}{s^2-4s+5} + \frac{2}{(s^2-4s+5)(s-2)^3}$.**Step 3:** Compute the inverse Laplace transform of $Y(s)$:

$$\begin{aligned} Y(s) &= \frac{4(s-2)}{(s-2)^2+1} + \frac{2}{(s-2)^2+1} + \frac{-2}{s-2} + \frac{2}{(s-2)^3} \\ \implies &\boxed{y(t) = 4e^{2t} \cos t + 2e^{2t} \sin t - 2e^{2t} + t^2 e^{2t}}. \end{aligned}$$

Partial fraction decomposition:

$$\frac{2s - 2}{s^2 - 4s + 5} = \frac{2(s - 2)}{(s - 2)^2 + 1} + \frac{2}{(s - 2)^2 + 1}$$

$$\frac{2}{(s^2 - 4s + 5)(s - 2)^3} = \frac{A(s - 2) + B}{(s - 2)^2 + 1} + \frac{C}{s - 2} + \frac{D}{(s - 2)^2} + \frac{E}{(s - 2)^3}$$

Tedious to calculate ...

Tip 1: Working in \mathbb{C} makes life easier!

$$\begin{aligned} \frac{2}{(s^2 - 4s + 5)(s - 2)^3} &= \frac{A(s - 2) + B}{(s - 2)^2 + 1} + \frac{C}{s - 2} + \frac{D}{(s - 2)^2} + \frac{E}{(s - 2)^3} \\ &= \frac{F}{s - 2 - i} + \frac{F^*}{s - 2 + i} + \frac{C}{s - 2} + \frac{D}{(s - 2)^2} + \frac{E}{(s - 2)^3} \end{aligned}$$

$$F = \left[\frac{2}{\cancel{(s - 2 - i)}(s - 2 + i)(s - 2)^3} \right]_{s=2+i} = \frac{2}{2i \cdot i^3} = 1$$

$$A = F + F^* = 2\operatorname{Re}\{F\} = 2,$$

$$B = i(F - F^*) = -2\operatorname{Im}\{F\} = 0$$

Tip 2: Taking derivatives

$$\frac{2}{(s^2 - 4s + 5)(s - 2)^3} = \frac{2(s - 2)}{(s - 2)^2 + 1} + \frac{C}{s - 2} + \frac{D}{(s - 2)^2} + \frac{E}{(s - 2)^3}$$

$$E = \left[\frac{2}{s^2 - 4s + 5} \right]_{s=2} = 2$$

$$D = \left[\frac{d}{d(s - 2)} \frac{2}{s^2 - 4s + 5} \right]_{s=2} = \left[\frac{-2}{[(s - 2)^2 + 1]^2} 2(s - 2) \right]_{s=2} = 0$$


$$\begin{aligned} C &= \left[\frac{1}{2!} \frac{d^2}{d(s - 2)^2} \frac{2}{s^2 - 4s + 5} \right]_{s=2} \\ &= -2 \left[\frac{[(s - 2)^2 + 1]^2 - 2(s - 2)^2 2[(s - 2)^2 + 1]}{[(s - 2)^2 + 1]^4} \right]_{s=2} = -2 \end{aligned}$$

Translation on the t -Axis


Let's compute $\mathcal{L}\{f(t-a)\}$, $a > 0$, given that $\mathcal{L}\{f(t)\} = F(s)$:

$$\begin{aligned}
 \mathcal{L}\{f(t-a)\} &= \int_0^{\infty} f(t-a)e^{-st} dt \stackrel{\tau:=t-a}{=} \int_{-a}^{\infty} f(\tau)e^{-s(\tau+a)} d\tau \\
 &= e^{-as} \int_0^{\infty} f(\tau)e^{-s\tau} d\tau + e^{-as} \int_{-a}^0 f(\tau)e^{-s\tau} d\tau \\
 &= e^{-as} F(s) + e^{-as} \int_{-a}^0 f(\tau)e^{-s\tau} d\tau \\
 &= \boxed{e^{-as} F(s)}, \quad \text{if } f(t) = 0 \text{ when } t < 0
 \end{aligned}$$

If $f(t) = 0$ for $t < 0$, then $\mathcal{L}\{f(t-a)\} = e^{-as}\mathcal{L}\{f(t)\}$, for $a > 0$.



How about functions $f(t)$ that is non-zero for $t < 0$?




Unit Step Function

Definition (Unit Step Function)

$$\mathcal{U}(t) := \begin{cases} 1, & t \geq 0 \\ 0, & t < 0 \end{cases}.$$


Note: $\mathcal{L}\{f(t)\mathcal{U}(t)\} = \mathcal{L}\{f(t)\}$.



Theorem (Translation on the t -Axis)

For $a > 0$,

$$\mathcal{L}\{f(t-a)\mathcal{U}(t-a)\} = e^{-as}\mathcal{L}\{f(t)\mathcal{U}(t)\} = e^{-as}\mathcal{L}\{f(t)\}.$$



Examples: Laplace Transforms

Example

Calculate $\mathcal{L}\{U(t-a)\}$.

$$\mathcal{L}\{U(t-a)\} = \mathcal{L}\{1 \cdot U(t-a)\} = e^{-as} \mathcal{L}\{1\} = \frac{e^{-as}}{s}.$$

Example

Calculate $\mathcal{L}\{\cos t U(t-\pi)\}$.

$$\begin{aligned} \mathcal{L}\{\cos t U(t-\pi)\} &= \mathcal{L}\{\cos(t+\pi-\pi)U(t-\pi)\} \\ &= e^{-\pi s} \mathcal{L}\{\cos(t+\pi)\} = e^{-\pi s} \mathcal{L}\{-\cos t\} \\ &= -e^{-\pi s} \frac{s}{s^2+1}. \end{aligned}$$

Examples: Inverse Laplace Transforms

Example

$$\text{Calculate } \mathcal{L}^{-1} \left\{ \frac{1}{(s-4)e^s} \right\}.$$

Example

$$\text{Calculate } \mathcal{L}^{-1} \left\{ \frac{se^{-2s} + e^{-s}}{(s-1)(s-2)} \right\}.$$