



## RESEARCH ARTICLE

## A Rule-Based Species Predictive Model for the Vulnerable Fairy Pitta (*Pitta nympha*) in Taiwan

Chia-Ying Ko<sup>(1)</sup>, Pei-Fen Lee<sup>(1)</sup>, Mei-Ling Bai<sup>(2)</sup> and Ruey-Shing Lin<sup>(3\*)</sup>

1. Institute of Ecology and Evolutionary Biology, National Taiwan University, 1, Roosevelt Rd., Sec. 4, Taipei 106, Taiwan; equal contribution.

2. Research Center for Biodiversity, Academia Sinica, 128, Sec. 2, Academia Rd., Taipei 115, Taiwan.

3. Endemic Species Research Institute, 1, Ming-Shen E. Road, Nantou 552, Taiwan.

\* Corresponding author. Tel: 866-49-2761331 ext. 142; Fax: 866-49-2761583; Email: rslin@tesri.gov.tw

(Manuscript received 9 July 2008; accepted 15 October 2008)

**ABSTRACT:** A fundamental step in biodiversity conservation is to identify potential distribution and quality habitat for a desired species, especially when the target is rare and difficult to detect. We demonstrated a GIS application in developing a quick predictive model to study the globally vulnerable Fairy Pitta (*Pitta nympha*) in Taiwan. We compiled sighting data between 1982 and 2000, established a rule-based model to predict its distribution, and applied the predictive map to design a sampling protocol, conducted field surveys to evaluate the accuracy of the model and to obtain hotspots. The results showed that most known distribution of the Fairy Pitta occurred in low elevation, hilly and forested areas. The map predicted 21.6% areas of Taiwan suitable for the Fairy Pitta and 78% of them occurred in western Taiwan. A total of 511 pittas were detected during the 2001 survey that covered 4% areas of Taiwan or 14.3% of predictive areas, with a mean of 30.2% detection probability per grid cell (2 × 2 km in resolution). The adjusted data indicated that the overall accuracy of our model was increased to 40.3% with 290 qualified cells. Most of the new sightings of the Fairy Pitta arising from the 2001 field survey fell in our predictive areas with most of them occurring in western Taiwan. The probability of detecting pitta was highest in the active selection cells within predictive areas. Based on mean number of pitta detected per cell, the hotspots of the Fairy Pitta in Taiwan included three regions: the watershed of Shimen Reservoir within Hsinchu and Taoyuan County, Linnei of Yulin County and the watershed of Wusanto Reservoir in Tainan County. We concluded that the model provides quick and effective predictions for planning conservation strategies and is particularly useful for rare species.

**KEY WORDS:** Niche, rule-based model, spatial sampling, vulnerable species, macrohabitat.

### INTRODUCTION

The Fairy Pitta (*Pitta nympha*) is currently listed as a vulnerable species by the BirdLife International Red Data Book (BirdLife International, 2001) and IUCN Red List of Threatened Species (IUCN, 2007), and its trading is also restricted by CITES (Erritzoe and Erritzoe, 1993; Inskipp and Gillett, 2005; CITES Species Database, 2008). Distributed in East Asia, the bird is migratory, and its worldwide population size may be less than 10,000 individuals (BirdLife International, 2001). However, information of Fairy Pitta worldwide is scattering and limited. Due to its distribution in the low-elevation regions that are highly susceptible to human activities, conservation efforts of the Fairy Pitta is urgent (Lambert and Woodcock, 1996; BirdLife International, 2001). In Taiwan, the Fairy Pitta is a summer visitor and previous studies (e.g., Severinghaus et al., 1991) have shown limited sighting records.

Geographic Information System (GIS) has become a widely-used tool to investigate species distribution (Smith-Ramirez et al., 2007; Isaac et al., 2008) and macrohabitat characteristics (Murgante and Las Casas, 2004; Uy and Nakagoshi, 2008). Digital

environmental information layers relating to a target species can be analyzed to identify potentially suitable habitats (Vogiatzakis, 2003). Conservation applications using GIS, for example species predictive models, have emerged in the last decade to help in guiding future survey efforts for rare and endangered species (Guisan and Zimmermann, 2000; Walther et al., 2007).

Designing a field sampling scheme, particularly for a rare species faces great challenges due to limited information. Spatial distribution modeling provides relatively accurate information when preparing for field survey. With this help, researchers can design better and efficient sampling plan to gather field information in a short time period (Powell et al., 2005; Walther et al., 2007) that would save time and money.

We hypothesized that the distribution of the vulnerable Fairy Pitta can be formulated using existing know distribution. In this study, we demonstrated the development of a predictive distribution model using GIS to study the Fairy Pitta in Taiwan. In particular, we compiled known sighting data between 1982 and 2000, identified macrohabitat characteristics using environmental GIS layers, and established a rule-based model to predict the potential distribution range of pitta. Finally, we applied the predictive map to design a field



sampling protocol and used these results to evaluate the accuracy of the model and to pinpoint hotspots of the species.

## MATERIALS AND METHODS

### Species data

We extracted and compiled sighting data between 1982 and 2000 from the databases of Wild Bird Federation Taiwan and Endemic Species Research Institute. Due to the elusive nature of the species, there were only 59 records. All sighting records were recorded between mid-April and September, the breeding season of the species, thus the sighting locations represented the breeding grounds of the species in Taiwan. These were the best available distribution of Fairy Pitta in Taiwan in 2000. Using a 2 × 2 km GIS grid representation (9,388 in total, Lee et al., 2004), we mapped the occurrence data into 54 cells (Fig. 1).

### Environmental data

To explore macrohabitat characteristics of the Fairy Pitta, we applied 13 environmental variables derived from a GIS database of ecological factors in Taiwan (Lee et al., 1997). The 13 variables were chosen by consulting previous literature (e.g., Severinghaus, 1991) that mentioned the Fairy Pitta ecology and our field experience. The GIS layers were also in the 2 × 2 km grid system. These variables can be classified into three categories, i.e., topography, climate and human activity (Table 1). We calculated mean and minimum elevation layers based on a digital elevation model (DEM, 40 × 40 m in resolution) produced by the Aerial Survey Office of Forest Bureau, Taiwan. We also calculated the coefficient of variation of elevation. Through the river and dam maps, we calculated the nearest distance from each cell

to the rivers. Climate variables included total annual precipitation and mean monthly temperature between 1959 and 1985. The 12 monthly total precipitation data were transformed into four PC variables (axes) by PCA (principal component analysis). The first PC variable, explaining 42% of the total variations in the 12 monthly total precipitation, represented characteristics of the precipitation between September and November. The second axis had the precipitation characteristics of May to September and explained 33% of the variations. The third (10%) variable characterized precipitation patterns from November to June, and the fourth (7%) variable represented precipitation of July. Warmth index reflected vegetation growth condition and was calculated by summing all the monthly mean temperature that was higher than 5°C (Liu and Su, 1992). Urbanization index and nearest distance to 3,000 m a.s.l. or above was the only two variables representing human activity in our study. Distance to 3,000 m a.s.l. or above was derived from the mean elevation layer. Urbanization index was calculated by combining the structure of employment populations from each township compiled from the Directorate General of Budget, Accounting and Statistics of Executive Yuan, in particular, the index was calculated as a sum of a second (mining, water, electricity, and building industry etc.) and a third classes (business, transportation, communication and service etc.) of industrial population divided by a sum of a first (agriculture), a second and a third classes of industrial population (see Sun et al., 1988 for detail calculations).

### Model construction

We first examined the relationships between the occurrence records and the environmental variables by performing an overlay analysis, and later an exploratory data analysis. Based on prior understanding of the Fairy Pitta, we assumed that all variables we chose were

**Table 1. Environmental variables used in this study.**

Category	Abbreviation	Variable description and unit
Topography	D2_RIVER	Nearest distance to main rivers and dams (km)
	ELEV	Mean elevation (m)
	ELEVMIN	Minimum elevation (m)
	ELEVCV	Coefficient of variation of elevation
Climate	P_TOTAL	Total annual precipitation (mm)
	P_PC1	The first PC axis (explaining 41.9% of the total variations) of 12 monthly total precipitation
	P_PC2	The second PC axis (33.0%) of 12 monthly total precipitation
	P_PC3	The third PC axis (10.2%) of 12 monthly total precipitation
	P_PC4	The fourth PC axis (6.5%) of 12 monthly total precipitation
	T_PC1	PC axis (explaining 96.5% of the total variations) of 12 monthly mean temperature
Human activity	WARMTH	Warmth index which sums mean monthly temperature higher than 5°C in a year (°C)
	URBAN	Urban development index
	D2_3000	Nearest distance to elevation 3,000 m and higher (km)



indicators of the bird's distribution in Taiwan. Some variables, such as elevation, might represent an upper limit to the distribution; some were a minimum requirement and some, such as temperature, showed an optimal range. We then determined the macrohabitat characteristics of the Fairy Pitta by examining the mean, standard deviation, minimum and maximum values of these variables through descriptive statistics. Based on the environmental characteristics of previous sighting data, we used the minimum and maximum values of these 13 environmental variables as the boundaries to develop 13 initial suitability map for the Fairy Pitta. Each map differentiated suitable and unsuitable habitat areas for the bird. These maps were then intersected to derive a map with predictive areas that met all the requirements. Since the birds mainly distributed in forests, our final predictive map excluded unforested areas by using a forested cover map.

### Model evaluation

In order to evaluate prediction accuracy of the predictive map, in March 2001 we designed a field sampling protocol based on the predictive map (Fig. 2). We identified three types of sampling areas from the map: (1) random cells within predictive areas, (2) active selection cells within predictive areas, and (3) active selection cells outside predictive areas. The random cells were randomly chosen within the predictive areas. The active selection cells were selected based on known sighting locations (Fig. 1) and our field experience. These active sites concentrated in the Shimen Reservoir watershed of Taoyuan County and the proposed Fushan Dam (Linnei) of Yunlin County. We also used our knowledge to choose the active selection cells outside predictive areas, although they might represent marginal habitats for pittas. The first two types were designed to evaluate the accuracy of predicting the presence of the Fairy Pitta, and the third one was helpful in checking the model's omission error. A total of 378 cells, including 208 random cells, 63 active selection cells within predictive areas, and 107 active selection cells outside predictive areas, were selected. Together, those cells covered 4% of Taiwan. Within each cell, we selected at least 3 sampling points to conduct the field survey. A total of 4,239 points, including 2,596 in random cells, 526 in active selection cells within predictive areas, and 1,117 in active selection cells outside predictive areas, were surveyed.

The field survey was conducted at each point from late April to early July in 2001. A total of more than 250 volunteer surveyors participated in the field survey. Each surveyor was trained with standardized

sampling protocol before the field surveys. For the surveys, we applied a playback-response method (Lin et al., 2007) to get quick responses of the Fairy Pitta. The method greatly helped in detecting the presence of birds (Lin et al., 2007).

We calculated the mean number of cells with pitta detected and the mean number of sampling point per cell in three sampling selection by using complete data. Since low sampling efforts, unequal spatial coverage, and incorrect survey timing might reduce the detection probability of Fairy Pitta (Lin et al., 2007), we further adjusted the survey data by deleting those cells with < 6 sampling points, less spatial coverage (< 50% of the areas) and surveyed outside the preferred survey period (late April to end of May) and calculated the same parameters.

We performed GIS operations in IDRISI and ArcGIS environments. All statistical analyses were conducted using SYSTAT and SAS.

## RESULTS

### Macrohabitat Preferences

Most known distributions of the Fairy Pitta occurred in forested lowland hills of western Taiwan (Fig. 1). Out of the 54 grid cells, only two sightings were located in the eastern Taiwan. The birds distributed islandwise but concentrated in the regions near Shimen Reservoir of Taoyuan and Hisichu County, and Tsengwen Reservoir of Chiayi County.

Comparing the sighting data with the environment ranges of the whole Taiwan, the Fairy Pitta preferred some environmental characteristics (Fig. 3 and Table 2). Since Fig. 3 showed the whole range value in Taiwan, the sighting records were mostly located in low elevation, moderate temperature, relatively high annual precipitation, and low urban development. In particular, the elevation ranges were between 0 and 980 m with 75% of the sightings occurred below 280 m a.s.l (Fig. 3a). The areas where the birds were found had higher temperature (Fig. 3b), and annual total precipitation ranged from 1,417 to 3,190 mm with a mean of 2,309 mm (Fig. 3c). These locations were also close to the rivers and dams (< 7.6 km), low urban development (Fig. 3d), and low warmth index.

### Predictive map

The predictive areas of the Fairy Pitta included 21.6% of the main island of Taiwan with 78% occurring in the western part of Taiwan (Fig. 2). Totally there were 2,031 cells predicted to have pittas. Overall, our prediction indicated the bird distributed in the forested region below 1,000 m a.s.l and away from high human



**Table 2. Summary of environmental characteristics of Fairy Pitta (*Pitta nympha*) sighting locations (n = 54).**

Variables	Mean ± SD	Minimum	Maximum
D2_3000	42.4 ± 11.8	24.7	94.7
D2_RIVER	1.8 ± 1.6	0	7.60
ELEV	268 ± 231.7	0	981
ELEVMIN	150 ± 149.2	0	650
ELEVCV	2.15 ± 1.15	0	5.67
P_TOTAL	2,308 ± 369	1,417	3,190
P_PC1	-81.4 ± 110.8	-266.4	205.5
P_PC2	96.8 ± 160.8	-226.7	506.1
P_PC3	-31.7 ± 103.2	-205.3	159.3
P_PC4	50.1 ± 102.6	-128.7	214.0
T_PC1	431 ± 62.5	235	540
WARMTH	200 ± 21.1	135	234
URBAN	52.8 ± 17.5	17	95

**Table 3. Field survey results of Fairy Pitta in 2001. The adjusted data were modified by deleting the cells with < 6 sampling points, less spatial coverage (< 50% of the cell area) and outside the preferred survey period (late April to end of May).**

Types of dataset and sampling	Total number of cells sampled	Number of cells with Pitta detected	Total sampling points	Number of sampling point per cell (mean ± SD)	Number of Pitta detected	Number of Pitta detected per sampling point (mean ± SD)
<b>Complete data</b>						
Random cells within the predictive regions	208	69 (33%)	2,596	12.7 ± 4.0	226	0.09 ± 0.20
Active cells within the predictive areas	63	28 (44%)	526	8.4 ± 7.5	237	0.31 ± 0.61
Active cells outside the predictive areas	107	17 (16%)	1,117	10.6 ± 4.9	48	0.06 ± 0.21
<b>Adjusted data</b>						
Random cells within the predictive regions	177	60 (34%)	2,376	13.4 ± 3.2	192	0.08 ± 0.17
Active cells within the predictive areas	29	23 (80%)	449	15.5 ± 5.0	223	0.43 ± 0.42
Active cells outside the predictive areas	84	13 (16%)	1,036	12.3 ± 3.3	40	0.04 ± 0.16

disturbed regions. In contrast to the very few sighting records in eastern Taiwan, our model predicted 22% of these areas were potential habitats for pitta. However, no area within Ilan County was predicted.

**Model evaluation**

A total of 511 pittas within 114 cells was detected during the 2001 survey (Fig. 4, Table 3), with a mean of 30.2% detection probability per cell.

Despite similar sampling efforts in the three sampling categories (Table 3), the active selection cells within predictive areas had significantly higher number of pitta detected per point than the random cells within predictive areas and the active selection cells outside predictive areas (H = 50.3, Kruskal-Wallis test, p < 0.001). The percentage of the number of cells with pitta detected was also the highest in the active selection cells within predictive areas (Table 3). In contrast, the active selection cells outside predictive

areas had the lowest percentage. Using sampling point as the basis, the mean detected individual per point was 0.12. The overall accuracy of our predictive model was 35.8%.

Analyses on the adjusted data indicated that the overall accuracy of our model was increased to 40.3% with 290 qualified cells (Table 3). The detection probability per cell also increased to 33.1%. The three sampling types showed significant difference in the number of pitta per point (H = 53.1, Kruskal-Wallis test, p < 0.001). The percentage of the number of cells with pitta detected and the number of pitta detected per point were the highest in the active selection cells within predictive areas, and similarly, the active selection cells outside predictive areas had the lowest values. However, due to the high number of sampling points, the mean number of pitta detected per sampling point in the adjusted data was significantly lower than that of all data (0.10 vs. 0.12, Mann-Whitney U test, p < 0.05).




Fig. 1. Known distribution of the Fairy Pitta (*Pitta nympha*) based on the sighting data compiled by Wild Bird Federation Taiwan and Taiwan Endemic Species Research Institute between 1982 and 2000.

The hotspots, based on the mean number of pitta detected per grid in the adjusted data, included three regions: the watershed of Shimen Reservoir within Hsinchu and Taoyuan Counties, Linnei of Yulin County and the watershed of Wusanto Reservoir in Tainan County. The detection probability reached 0.9. These sites also had higher complexity of topography and secondary forested cover.

## DISCUSSION

Conservation of rare and endangered species often faces urgent challenge to gather sufficient data for policy making in a short time period. In this study we demonstrated a simple and quick approach to obtain adequate distribution data for a globally vulnerable species, the Fairy Pitta. By integrating sighting data from all possible sources, we developed a species predictive model and designed a standardized sampling protocol to evaluate the model effectiveness. The rule-based model applied niche concept and assumed all the potential variables had equal






Fig. 2. Predictive distribution of the Fairy Pitta (*Pitta nympha*) and field sampling design. Three categories of sampling cells were included: i.e., random cells within: random cells within the predictive areas, active selection cells within: active selection cells within the predictive areas, and active selection cells outside: active selection cells outside the predictive areas. All the selections were based on the predictive areas and were used to conduct field surveys.

contributions. The resulting 40% overall predictive accuracy indicated that our model and the overall approach was straightforward and valuable. We also conducted similar field work in 2002 and 2005, the results suggested that the model accuracy could reach over 50% (R. S. Lin, unpublished data). In addition, the gathered information provides sufficient pitta presence data for decision makers and further insight on the model refinements (Lee et al., 2006).

We further applied the sampling results (i.e., Fig. 4) to develop more sophisticated models using logistic regression, discriminant function analysis, GARP, fuzzy logic and artificial neural network using a  $1 \times 1$  Km grid dataset. Instead of treating all explanatory variables as equal, these new models tried to weight the variables differently. Our results indicated that overall accuracies ranged from 63% to 85% and the GARP model had the highest predictive power (Lee et al., 2006). With the



**Fig. 3.** Overall ranges of environmental characteristics in Taiwan (in open bar) and macrohabitat characteristics of Fairy Pitta (*Pitta nympha*) (in solid bar) based on the sighting data compiled by Wild Bird Federation Taiwan and Taiwan Endemic Species Research Institute between 1982 and 2000: (a) mean elevation (m), (b) mean temperature (°C), (c) total annual precipitation (mm), and (d) urban development index. The data clearly indicated that the distribution of Fairy Pitta was restricted to some specific range.



**Fig. 4.** Presence data of the Fairy Pitta (*Pitta nympha*) in Taiwan after the 2001 survey that covered 4% in Taiwan or 14.3% in predictive areas.

ensemble forecasting approach (Araújo and New, 2007), we further developed a robust predictive model for Fairy Pitta. The combined model yielded a narrower predictive range than the rule-based model in our study, though the overall pattern was similar.

A vulnerable species is usually characterized with low population and easily threatened by extrinsic factors (Smith et al., 2003; Mitchell and Wilson, 2007). In the Fairy Pitta case, its elusive nature leads to difficulties in detecting presence in the field (Severinghaus et al., 1991; Lin et al., 2007). Their occurrence records are usually too few to build strong predictive models. Our rule-based model is simple to construct once the environmental dataset is available. As we demonstrated, the model acted as the initial guidance to design a systematic field sampling strategy. With the help of the playback-response method (Lin et al., 2007) and more than 250 field surveyors, the results of the field work provided more distribution data (108 cells) than those accumulated for almost 20 years (54 cells). Although these research efforts were different, our approach indicated that a systematic sampling design based on this rule-based model prediction substantially improved our knowledge on the distribution of the Fairy Pitta (Lin et al., 2007).

The number of pitta detected per sampling point or the number of pitta detected per cell in the active



selection cells outside predictive region was significantly lower than those in the random cells and active selection cells within predictive regions. Furthermore, within the predictive areas, the results that the active selection cells had far better detection probabilities than the random cells also indicated the effectiveness of the predictive model. These results clearly suggested the model provided better predictive power than random selection. Finally, species abundance distribution can be applied to integrate the single species prediction and environmental indicators (McGill et al., 2007).

As expected, the current model did not predict all possible occurrence areas of the Fairy Pitta in Taiwan. The number of pitta detected per cell in the active selection cells outside predictive areas clearly indicated that the potential omission error in the model. In particular, the Ilan County was completely missing in the prediction. Part of the reasons for the error was the few sighting records available to frame our selection rules. In contrast, some areas predicted by the current model, though not verified in the 2001 survey, were found to have pitta presences in either 2002 or 2005 surveys or in some scattered news reports (L. T. Chao, personal communication).

Although there are some advocacy in promoting the further applications of predictive model (e.g., Rodriguez et al., 2007), most species predictive mappings stopped after the prediction models were compared and the best one singled out (e.g., Maynard and Quinn, 2007). To our knowledge, our efforts represented a successful application to fully use the predictive model to obtain ecological distribution data on the Fairy Pitta in its breeding range. Without further affirmation, both predictive model and mapping cast doubts about their accuracy and usability (Loehle and LeBlanc, 1996; Travis, 1996). In our study, we showed the applicability of the model by extensive field work, and the probability to detect the Fairy Pitta was higher within predictive areas than non-predictive areas.

Given the results of our analyses, we proposed that it is critical and important to design an immediate conservation strategy for this vulnerable species. According to the model prediction (Fig. 4), the Fairy Pitta widely distributed at low elevation of the evergreen broad-leaved forest in Taiwan. Further gap analysis using all protected areas in Taiwan, including national parks, nature reserves, wildlife refuges and important wildlife habitats, indicated that only less than 2% of the currently known distributions of the Fairy Pitta were protected (Lee et al., 2006). Most of the distributions are located in the relatively high human disturbance zone and within the private lands. These lands suffer excessive developments and have

numerous predators, such as snakes, mammals, birds and ants (Chen, 2007). Clearly these situations complicate the conservation of this species.

The current hotspots for the Fairy Pitta are located in the low elevation forests of highly complex topography. We identified three regions based on the number of pitta detected per cell. All sites were close to rivers or dams. Since the watersheds of dams are protected by laws, it is possible that Fairy Pitta habitats in these regions can receive better protection. Since our sampling efforts were not complete, we suspected there are many more similar unidentified hotspots in Taiwan. To effectively protect the species, we suggested land managers avoid planning large-scale development projects in similar habitats.

The rule-based model and sampling protocol developed in this study can be applied to conduct survey in other countries where the Fairy Pitta breeds. Since the playback-response approach is an effective method for surveying the species (see Lin et al., 2007), with the abundant spatial and land cover data already available (Global Land Cover Facility, 2008), the only modification of the methodology may be the survey period because the birds breed at other latitudes have different timing comparing to those in Taiwan.

## ACKNOWLEDGEMENTS

We thank Endemic Species Research Institute of Council of Agriculture, Central Water Resources of the Ministry of Economic Affairs, and National Science Council for providing research funding and Council of Agriculture for support in environmental database construction. The help from more than 250 bird watchers in Taiwan greatly enhanced our knowledge on the Fairy Pitta. The comments by two anonymous reviewers greatly improved the manuscript.

## LITERATURE CITED

- Araújo, M. B. and M. New.** 2007. Ensemble forecasting of species distributions. *Trends in Eco. & Evol.* **22**: 42-47
- BirdLife International.** 2001. Threatened birds of Asia: the BirdLife International Red Data Book. BirdLife International, Cambridge, UK. 852pp.
- Chen, W.-J.** 2007. Predation risk of the shrub and ground nesting passerines in a disturbed lowland forest of central Taiwan. M. S. Thesis, National Taiwan University, Taipei, Taiwan. 68pp. (In Chinese, English Abstract)
- CITES Species Database.** 2008. CITES Species Database. Available at <http://www.cites.org/eng/resources/species.html>.
- Erritzoe, J. and H. B. Erritzoe.** 1993. The birds of CITES and how to identify them. Lutterworth Press, Cambridge, Great Britain, UK. 224pp.
- Global Land Cover Facility.** 2009. Global Land Cover Facility. Available at <http://www.landcover.org/index.shtml>.



- Guisan, A. and N. E. Zimmermann.** 2000. Predictive habitat distribution models in ecology. *Ecol. Mod.* **135**: 147-186.
- Inskipp, T. and H. J. Gillett, editors.** 2005. Checklist of CITES species and Annotated CITES Appendices and reservations. Compiled by UNEPWCMC. CITES Secretariat, Geneva, Switzerland and UNEP-WCMC, Cambridge, UK. 339 pp. & CD-ROM.
- Isaac, B., R. Cooke, D. Simmons and F. Hogan.** 2008. Predictive mapping of powerful owl (*Ninox strenua*) breeding sites using Geographical Information Systems (GIS) in urban Melbourne, Australia. *Landsc. Urban Plan.* **84**: 212-218.
- IUCN.** 2008. IUCN Red list of threatened species. Available at <http://www.iucnredlist.org>.
- Lambert, F. and M. Woodcock.** 1996. Pittas, broadbills and asities. Pica, Sussex, UK. 181pp.
- Lee, P.-F., C.-Y. Liao, T.-C. Lee, Y.-H. Pan, W.-H. Fu and H.-W. Chen.** 1997. An ecological and environmental GIS database for Taiwan. National Taiwan University, Taiwan. 105pp. (In Chinese, English abstract).
- Lee, P.-F., M.-L. Bai and R.-S. Lin.** 2006. Habitat preference and distribution prediction of vulnerable Fairy Pitta (*Pitta nympha*) in Taiwan by remote sensing and GIS. 2005 Scientific Progress in Remote Sensing Projects, Council of Agriculture, Taipei. pp. 1-10. (In Chinese, English abstract)
- Lee, P.-F., T.-S. Ding, F.-H. Hsu and S. Geng.** 2004. Breeding bird species richness in Taiwan: distribution on gradients of elevation, primary productivity and urbanization. *J. Biogeog.* **31**: 307-314.
- Lin, R.-S., P.-F. Lee, T.-Z. Ding and K. Y.-T. Lin.** 2007. Effectiveness of playbacks in censusing the Fairy Pitta (*Pitta nympha*) during the breeding season in Taiwan. *Zool. Stud.* **46**: 242-248.
- Liu, T.-R. and H.-J. Su.** 1992. Forest plant ecology. The Commercial Press, Ltd, Taipei, Taiwan. 490pp. (In Chinese)
- Loehle, C. and D. LeBlanc.** 1996. Model-based assessments of climate change effects on forests: A critical review. *Ecol. Model.* **90**: 1-31.
- Maynard, C. N. and J. F. Quinn.** 2007. Predicting species distribution: a critical comparison of the most common statistical models using artificial species. *J. Biogeog.* **34**: 1455-1469.
- McGill, B. J., R. S. Etienne, J. S. Gray, D. Alonso, M. J. Anderson, H. K. Benecha, M. Dornelas, B. J. Enquist, J. L. Green, F. He, A. H. Hurlbert, A. E. Magurran, P. A. Marquet, B. A. Maurer, A. Ostling, C. U. Soykan, K. I. Ugland and E. P. White.** 2007. Species abundance distributions: moving beyond single prediction theories to integration within an ecological framework. *Ecol. Lett.* **10**: 995-1015.
- Mitchell, E. and B. A. Wilson.** 2007. Detection and habitat use of the Rufous Bristlebird (*Dasyornis broadbenti*) in coastal heathland, in south-western Victoria, Australia. *EMU* **107**: 327-334.
- Murgante, B. and G. Las Casas.** 2004. GIS and fuzzy sets for the land suitability analysis. *Computational Science and Its Applications - ICCSA 2004, PT 2*, 3044, 1036-1045.
- Powell, M., A. Accad and A. Shapcott.** 2005. Geographic information system (GIS) predictions of past, present habitat distribution and areas for re-introduction of the endangered subtropical rainforest shrub *Triunia robusta* (Proteaceae) from south-east Queensland, Australia. *Biol. Conserv.* **123**: 165-175.
- Rodríguez, J. P., L. Brotons, J. Bustamante and J. Seoane.** 2007. The application of predictive modeling of species distribution to biodiversity conservation. *Diversity and Distrib.* **13**: 243-251.
- Severinghaus, L. L., C.-T. Liang, S. R. Severinghaus and L.-C. Lo.** 1991. The distribution, status and breeding of Fairy Pitta (*Pitta nympha*) in Taiwan. *Bull. Inst. Zool. Acad. Sin.* **30**: 41-47.
- Smith, S. J., Z. Lucas and W. T. Stobo.** 2003. Estimate of the Ipswich Sparrow population on Sable Island, Nova Scotia, in 1998, using a random-transect survey design. *Can. J. Zool.* **81**: 771-779.
- Smith-Ramirez, C., I. Diaz, P. Pliscoff, C. Valdovinos, M. A. Mendez, J. Larrain and H. Samaniego.** 2007. Distribution patterns of flora and fauna in southern Chilean coastal rain forests: Integrating natural history and GIS. *Biodiver. & Conser.* **16**: 2627-2648.
- Sun, C.-H., C.-Y. Chang and C.-L. Chang.** 1988. Research on drawing main features and establishing information system in Taiwan. National Taiwan University, Taipei, Taiwan. pp. 85-102. (In Chinese, English abstract).
- Travis, J.** 1996. The significance of geographical variation in species interactions. *Amer. Nat.* **148**: S1-S8.
- Uy, P. D. and N. Nakagoshi.** 2008. Application of land suitability analysis and landscape ecology to urban green space planning in Hanoi, Vietnam. *Urban For. & Urban Greening* **7**: 25-40.
- Vogiatzakis, I. N.** 2003. GIS-based modeling and ecology: a review of tools and methods. Geographical Paper No. 170. University of Reading, Whiteknights, UK. 34pp.
- Walther, B. A., N. Schaffer, A. van Niekerk, W. Thuiller, C. Rahbek and S. L. Chown.** 2007. Modeling the winter distribution of a rare and endangered migrant, the Aquatic Warbler *Acrocephalus paludicola*. *Ibis* **149**: 701-714.





## 臺灣八色鳥(*Pitta nympha*)之規則化分布預測模式

柯佳吟<sup>(1)</sup>、李培芬<sup>(1)</sup>、白梅玲<sup>(2)</sup>、林瑞興<sup>(3\*)</sup>

1. 國立臺灣大學生態學與演化生物學研究所，106 臺北市羅斯福路 4 段 1 號，臺灣。

2. 中央研究院生物多樣性研究中心，115 台北市南港區研究院路 2 段 128 號，臺灣

3. 行政院農業委員會特有生物研究保育中心，552 南投縣集集鎮民生東路 1 號，臺灣。

\* 通信作者。Tel: 866-49-2761331 ext. 142; Fax: 886-49-2761583; Email: rslin@tesri.gov.tw

(收稿日期：2008 年 7 月 9 日；接受日期：2008 年 10 月 15 日)

摘要：確認稀有性物種的分布和其適宜性棲地是生物多樣性保育的基本工作。本研究展現利用地理資訊系統，針對臺灣的全球易危性物種——八色鳥 (*Pitta nympha*)，發展一個簡易又快速的分布預測模式。收集 1982 至 2000 年的觀察記錄，建立八色鳥的規則化分布預測模式，並利用預測圖設計野外調查，以標準化的方法評估此模式的效能，建立熱點位置。結果顯示，八色鳥分布於低海拔且有森林覆蓋的丘陵地區。預測圖估計台灣約有 21.6% 的區域，可能會有八色鳥的出現，其中約有 78% 處在西部。在 2001 年的調查（調查區域約佔全台 4%，或預測區的 14.3%）中，共發現 511 隻八色鳥，平均每一個 2 × 2 公里網格的發現機率為 30.2%，以比較嚴格篩選的網格資料（共 290 格），模式準確度則為 40.3%。新發現有八色鳥出現的網格分布，大致與預測模式的型態相似，且 93.9% 出現在臺灣西部。每網格內的發現機率，以在預測區域內的刻意選擇區最高。以網格的出現機率為標準，八色鳥熱點出現在新竹與桃園縣石門水庫附近、雲林縣的林內鄉、臺南縣烏山頭水庫附近的山區。本模式特別適合稀有性物種，方法簡易又有效，可以協助瞭解物種的可能分布範圍，進而提供更適宜的保育策略。

關鍵詞：生態棲位、規則化模式、空間取樣、易危物種、巨觀棲地。