

Notes on Calabi-Yau Ordinary Differential Equations^{*†}

JENG-DAW YU[‡]

June 26, 2009

Abstract

We investigate the structures of Calabi-Yau differential equations and the relations to the arithmetic of the pencils of Calabi-Yau varieties behind the equations. This provides explanations of some observations and computations in the recent paper [6].

0 Introduction

This note may be regarded as a supplement to the paper [6] of Samol and van Straten. In that paper, the authors study the variation, along a projective line, of the Frobenius action on the cohomology of a Calabi-Yau variety (over a finite field) via the expansion at a totally degenerate point of a certain period of (a lifting of) the pencil. For certain families, they provide a p -adic analytic formula for the unit root as well as the Frobenius polynomial (in the case of rank 4) of each fiber provided the fiber is ordinary. Their explicit computations for those examples also reveal the existence of Dwork type congruences among the coefficients of the period of such a family.

In this note, we provide some theoretical explanations of the observations and computations in their paper. In particular, we show that certain cases of the Dwork congruences, including the mod p ones, follow from the relative geometry of the family.

This article is organized as follows. In §1, we explore the notion of ordinary differential equations of Calabi-Yau type over a field of characteristic 0 and derive some basic algebraic properties of them. This part provides a further investigation of abstract Calabi-Yau differential equations. In §2, families of Calabi-Yau varieties over a projective line with totally degenerate fibers at the origin are introduced. The associated Picard-Fuchs equations of such families provide the geometric source of the Calabi-Yau differential equations. §3 is devoted to the study of the mod p and p -adic properties of the Calabi-Yau equations. In particular, we study the relation between the solutions of a Calabi-Yau differential equation and the p -adic Hasse invariant of the underlying family. This relation provides an explanation of parts of the Dwork congruences. Unfortunately at this stage, we do not know how to deal with these Dwork congruences in general. Finally a further inspection of the Hadamard products appeared in [6] is given in the appendix.

*2010 MSC: 14J32, 12H05

†This work was partially supported by the 2008 Starting Grant of College of Science, NTU, Grant 97-2115-M-002-018 of NSC, and a grant of NTU, Taiwan.

‡Department of Mathematics, National Taiwan University, Taipei 10617, Taiwan; E-mail address: jdyu@math.ntu.edu.tw

1 Ordinary differential equations of Calabi-Yau type

This section consists of exercises in the theory of differential modules. For basic references, see [8, Chap 2]. We recall the notion of differential equations of Calabi-Yau type and derive some basic properties of them. Such equations arise in the study of the Picard-Fuchs equations of pencils of Calabi-Yau varieties with certain degenerations. We postpone the geometric picture to the following sections. Notice that some properties we derived here (e.g., Lemma 1.3) can be obtained more easily if the differential equations are from geometry.

Let K be a field of characteristic 0 with a fixed embedding into \mathbb{C} . Denote by $\overline{K} \subset \mathbb{C}$ the algebraic closure of K . Let t denote a variable. It will also be regarded as a fixed parameter of the projective line \mathbb{P}^1 over K . Let $\theta = t \frac{d}{dt}$ be the usual logarithmic derivation with respect to t . In this paper, we will use the highly *non-standard* convention:

$$g' := \theta g$$

for a differentiable function $g = g(t)$ of t throughout the discussion. Now consider an ordinary linear differential operator of order $n \geq 1$ of the form

$$\mathcal{L} = \theta^n + \sum_{i=0}^{n-1} a_i \theta^i \tag{1}$$

with coefficients $a_i = a_i(t) \in K(t)$.

(a) *The condition (N) and the β -factor*

Suppose \mathcal{L} is a differential operator of order n of the form (1). Consider the following condition on \mathcal{L} :

(N) Null exponents: \mathcal{L} has a regular singularity at the origin $t = 0$ (i.e., $a_i \in K(t) \cap K[[t]]$) and the associated indicial polynomial of \mathcal{L} at this point reduces to

$$s^n + \sum_{i=0}^{n-1} a_i(0) s^i = s^n. \tag{2}$$

For any \mathcal{L} as in (1), let formally

$$\beta := \exp \left(\frac{2}{n} \int a_{n-1} \frac{dt}{t} \right) \tag{3}$$

(i.e. a *non-zero* solution of $n\beta' = 2a_{n-1}\beta$). We call it the β -factor of \mathcal{L} . It exists at least in some differential field extension of $\overline{K}(t)$.

Lemma 1.1 *Suppose \mathcal{L} as in (1) satisfies the condition (N). Then we have the following.*

(i) *The β -factor in (3) can be taken from $(1 + tK[[t]])$.*

(ii) Regard \mathcal{L} as a differential operator in $\overline{K}(t)[\theta]$. Then the differential Galois group of \mathcal{L} is contained in $\mathrm{SL}(n)_{\overline{K}}$ if and only if β can be chosen with

$$\sqrt{\beta^n} \in K(t) \cap (1 + tK[[t]]).$$

Proof. By (2), $a_{n-1}(0) = 0$ and hence one can formally choose

$$\int a_{n-1} \frac{dt}{t} \in tK[[t]].$$

Thus the first assertion follows. The differential Galois group of $\mathcal{L} \in \overline{K}(t)[\theta]$ is in $\mathrm{SL}(n)$ if and only if there is a non-zero solution in $\overline{K}(t)$ of the operator $\theta + a_{n-1}$ ([8, Exercise 1.35.5]). Thus one can require that

$$\beta \in \sqrt[n]{\overline{K}(t)} \cap (1 + tK[[t]])$$

and (ii) follows accordingly. \square

From now on, we will always assume that $\beta \in (1 + tK[[t]])$ if \mathcal{L} satisfies the condition (N).

(b) *The self-adjointness and the polarization*

Let \mathcal{L} be as in (1). Recall that the *formal adjoint* \mathcal{L}^* of \mathcal{L} is the differential operator

$$\mathcal{L}^* = (-1)^n \theta^n + \sum_{i=0}^{n-1} (-1)^i \theta^i a_i.$$

We say that \mathcal{L} is *self-adjoint* if, as elements in $K(t)[\theta]$,

$$\mathcal{L}^* = (-1)^n \beta \mathcal{L} \beta^{-1}, \tag{4}$$

where β is defined in (3).

For any \mathcal{L} of order n , denote by $\mathcal{M}_{\mathcal{L}}$ the left $K(t)[\theta]$ -module with a generator η defined by:

$$\begin{aligned} K(t) &\rightarrow \mathcal{M}_{\mathcal{L}} := K(t)[\theta]/K(t)[\theta]\mathcal{L} \\ 1 &\mapsto \eta, \end{aligned} \tag{5}$$

where the map is the natural projection. As a $K(t)$ -module, $\mathcal{M}_{\mathcal{L}}$ is free of rank n with a basis $\{\eta^{(i)}\}_{i=0}^{n-1}$, where $\eta^{(i)} := \theta^i \eta$. An element $x \in \mathcal{M}_{\mathcal{L}}$ is called *horizontal* if $x' := \theta x = 0$.

Define a filtration Fil^\bullet on $\mathcal{M}_{\mathcal{L}}$ by setting Fil^i = the $K(t)$ -submodule generated by $\{\eta^{(j)}\}_{j=0}^{n-1-i}$. A *polarization* on $\mathcal{M}_{\mathcal{L}}$ is a $K(t)$ -linear, $(-1)^{n+1}$ -symmetric, non-degenerate horizontal pairing

$$\langle \cdot, \cdot \rangle : \mathcal{M}_{\mathcal{L}} \times \mathcal{M}_{\mathcal{L}} \rightarrow K(t)$$

such that $\langle \mathrm{Fil}^i, \mathrm{Fil}^{n-i} \rangle = 0$ for $0 \leq i \leq (n-1)$. As usual, we say $\mathcal{M}_{\mathcal{L}}$ is *polarizable* if there exists a polarization on it; $\mathcal{M}_{\mathcal{L}}$ is called *polarized* if it is equipped with an underlying polarization. The aim of this subsection is to prove the following.

Theorem 1.2 *Let \mathcal{L} be as in (1) and β be defined in (3). Then \mathcal{L} is self-adjoint with $\beta \in K(t)$ if and only if $\mathcal{M}_{\mathcal{L}}$ is polarizable.*

We first prove the following.

Lemma 1.3 *Let $\langle \cdot, \cdot \rangle$ be a $K(t)$ -linear horizontal pairing on $\mathcal{M}_{\mathcal{L}}$ such that $\langle \eta, \eta^{(i)} \rangle = 0$ for $0 \leq i < (n-1)$. Let $\gamma = \langle \eta, \eta^{(n-1)} \rangle$. We have*

- (i) *The pairing is uniquely determined by γ and $\gamma = c\beta^{-1}$ for some $c \in K$.*
- (ii) *The pairing is $(-1)^{n+1}$ -symmetric.*
- (iii) *The pairing is a polarization if $\gamma \neq 0$.*

Proof. Since $\langle \cdot, \cdot \rangle$ is horizontal, the values $\langle \eta, \eta^{(i)} \rangle, 0 \leq i \leq m$ uniquely determine $\langle \eta^{(i)}, \eta^{(j)} \rangle$ for all $i + j \leq m$ by a simple inductive procedure of taking derivatives. Thus in our case, the pairing is uniquely determined by γ . We have $\langle \text{Fil}^i, \text{Fil}^{n-i} \rangle = 0$ for $0 \leq i \leq (n-1)$ and the pairing is trivial if $\gamma = 0$.

Since $\langle \eta, \eta^{(n-2)} \rangle = 0$, we have

$$\begin{aligned} 0 &= \langle \eta, \eta^{(n-2)} \rangle' \\ &= \langle \eta', \eta^{(n-2)} \rangle + \langle \eta, \eta^{(n-1)} \rangle, \end{aligned}$$

which implies that $\langle \eta', \eta^{(n-2)} \rangle = -\gamma$. Similarly one finds, for $0 \leq i \leq n-1$, that

$$\langle \eta^{(i)}, \eta^{(n-1-i)} \rangle = (-1)^i \gamma$$

by induction. In particular, the pairing is non-degenerate if $\gamma \neq 0$.

Assume $n = 2l$ is even. Then

$$\begin{aligned} \gamma &= \langle \eta, \eta^{(n-1)} \rangle \\ &= -\langle \eta', \eta^{(n-2)} \rangle \\ &\quad \vdots \\ &= (-1)^{l-1} \langle \eta^{(l-1)}, \eta^{(l)} \rangle. \end{aligned}$$

Taking derivatives of the l equations above, rewriting $\eta^{(n)}$ in terms of $\{\eta^{(i)}\}_{i=0}^{n-1}$ via \mathcal{L} and summing them up, one gets

$$l\gamma' = -a_{n-1}\gamma.$$

Thus by definition, $\gamma = c\beta^{-1}$ for some $c \in K$. The computation is similar for n odd.

We now prove the parity of $\langle \cdot, \cdot \rangle$. We may assume that the pairing is non-degenerate. Let $\mathcal{N} = \text{Hom}_{K(t)}(\mathcal{M}_{\mathcal{L}}, K(t))$ and consider the natural pairing

$$(\cdot, \cdot) : \mathcal{N} \times \mathcal{M}_{\mathcal{L}} \rightarrow K(t)$$

given by $(f, m) = f(m)$. We equip \mathcal{N} with the differential module structure such that (\cdot, \cdot) is horizontal. Then indeed, as a $K(t)[\theta]$ -module, \mathcal{N} is generated by ξ with

$$(\xi, \eta^{(i)}) = \begin{cases} 1 & \text{if } i = n-1 \\ 0 & \text{otherwise} \end{cases}$$

([8, Exercise 2.12.6]). Denote $\xi^{(i)} = \theta^i \xi$. By a similar computation as above, one deduces that

$$(\xi^{(i)}, \eta^{(j)}) = \begin{cases} (-1)^i & \text{if } i + j = n - 1 \\ 0 & \text{otherwise.} \end{cases}$$

Now the non-degenerate pairing $\langle \cdot, \cdot \rangle$ induces an isomorphism f between $\mathcal{M}_{\mathcal{L}}$ and \mathcal{N} and it sends η to $\gamma \xi$. Thus under f , we can regard $\{\xi^{(i)}\}_{i=0}^{n-1}$ as another basis of $\mathcal{M}_{\mathcal{L}}$. Moreover, since \mathcal{N} and $\mathcal{M}_{\mathcal{L}}$ are dual to each other (indeed, $\mathcal{N} = \mathcal{M}_{\mathcal{L}^*}$), after switching the roles of $\mathcal{M}_{\mathcal{L}}$ and \mathcal{N} , we see that

$$\langle \xi^{(i)}, \eta^{(j)} \rangle = (-1)^{n-1} \langle \eta^{(j)}, \xi^{(i)} \rangle.$$

This completes the proof of the parity of $\langle \cdot, \cdot \rangle$. \square

Proof of Thm 1.2. Suppose \mathcal{L} is self-adjoint and $\beta \in K(t)$. Then by the Lemma above, we can equip $\mathcal{M}_{\mathcal{L}}$ with the polarization determined by setting $\langle \eta, \eta^{(n-1)} \rangle = \beta^{-1}$.

On the other hand, suppose $\mathcal{M}_{\mathcal{L}}$ is polarized. Then $\beta \in K(t)$ by the Lemma above. Multiplying by a non-zero constant, we may assume that $\langle \eta, \eta^{(n-1)} \rangle = \beta^{-1}$. Then $\beta \eta$ is dual to $\eta^{(n-1)}$ with respect to the basis $\{\eta^{(i)}\}_{i=0}^{n-1}$. Thus $\mathcal{L}^* \beta \eta = 0$ ([8, Exercise 2.12.6]) and hence $\mathcal{L}^* = (-1)^n \beta \mathcal{L} \beta^{-1}$. \square

(c) Calabi-Yau differential equations and the q -coordinate

Definition. A differential operator $\mathcal{L} \in K(t)[\theta]$ of the form (1) is called *Calabi-Yau* if \mathcal{L} is self-adjoint and satisfies condition (N) in §1(a).

Remark. It might be better to call such an \mathcal{L} as above *locally* or *quasi-Calabi-Yau* since in literature (e.g., in [1], [6]), there are some integral conditions on solutions of \mathcal{L} (cf. Thm 3.3) and here we do not require that \mathcal{L} is regular singular away from 0. However there seems to be no unified definition yet.

For any \mathcal{L} as in (1) satisfying the condition (N), we set

$$\Lambda_{\mathcal{L}} = K[[t]][\theta] / K[[t]][\theta] \mathcal{L}. \quad (6)$$

It is a $K[[t]]$ -lattice in the completion $\mathcal{M}_{\mathcal{L}} \otimes_{K[t]} K[[t]]$ of $\mathcal{M}_{\mathcal{L}}$. We abuse the notation by denoting $\text{Fil}^i \subset \Lambda_{\mathcal{L}}$ the induced filtration from $\text{Fil}^i \subset \mathcal{M}_{\mathcal{L}}$.

Suppose \mathcal{L} is Calabi-Yau. Denote by $F(t) = F_0(t)$ the unique formal power series such that

$$\mathcal{L}F(t) = 0, \quad F(t) \in 1 + tK[[t]]; \quad (7)$$

for $n \geq 2$, denote by $F_i(t), 1 \leq i \leq n-1$, the solutions with logarithmic pole of the form

$$\mathcal{L}F_i(t) = 0, \quad \sum_{r=0}^i (-1)^r F_{i-r}(t) \cdot \frac{\log^r t}{r!} \in tK[[t]]. \quad (8)$$

Thus $\{F_i\}$ forms the Frobenius basis of solutions near $t = 0$ of \mathcal{L} . Let

$$wr_i = wr(F, F_1, \dots, F_i) := \det \left(F_r^{(s)} \right)_{0 \leq r, s \leq i} \quad (9)$$

be the wronskians of $\{F_r\}_{r=0}^i$ and set

$$q = \exp\left(\frac{F_1}{F}\right) \in t + t^2 K[[t]]. \quad (10)$$

Thus $K[[t]] = K[[q]]$. We call q the q -coordinate of \mathcal{L} .

Theorem 1.4 *Suppose \mathcal{L} in (1) is Calabi-Yau. With the notations above, there exists a unique increasing filtration U_\bullet of $K[[t]][\theta]$ -submodules of $\Lambda_{\mathcal{L}}$ such that, for all i ,*

$$\Lambda_{\mathcal{L}} = U_i \oplus \text{Fil}^{i+1} \quad (11)$$

and U_{i+1}/U_i are trivial $K[[t]][\theta]$ -modules. Moreover, up to a multiplicative constant, there exists a unique sequence $\{u_i \in \Lambda_{\mathcal{L}}\}_{i=0}^{n-1}$ with the following two properties:

- (i) As a $K[[t]]$ -module, U_i is generated by $\{u_r\}_{r=0}^i$.
- (ii) u_i is of the form $u_i = \sum_{r=0}^{n-1-i} v_{i,r} \eta^{(r)}$ with

$$v_{i,n-1-i} = \beta \frac{wr_i}{wr_{i-1}}. \quad (12)$$

Consequently, we have $u'_{i+1} = \tau_{i+1} \cdot u_i$, where

$$\tau_{i+1} = \frac{wr_{i-1}wr_{i+1}}{wr_i^2},$$

and in particular, $\tau_1 = (\log q)'$.

Proof. We find $\{u_i\}$ satisfying (11) and (12) by induction.

The condition (N) of \mathcal{L} implies that, up to a constant multiple, there is a unique non-zero element

$$u_0 = \sum_{r=0}^{n-1} v_{0,r} \eta^{(r)}, \quad v_{0,r} \in K[[t]]$$

which is horizontal. Since

$$\mathcal{L}\langle u_0, \eta \rangle = \langle u_0, \mathcal{L}\eta \rangle = 0,$$

we see that $v_{0,n-1} = c\beta F$ for some constant c by Lemma 1.3. It is obvious that $c \neq 0$. Thus after modifying by a constant, $v_{0,n-1}$ is of the form (12).

Let $\mathcal{L}^{[i]}$ be the i -th exterior product of \mathcal{L} . Then $\mathcal{L}^{[i]}$ satisfies the condition (N) and $\Lambda_{\mathcal{L}^{[i]}}$ is a quotient of the i -th exterior power of the $K[[t]]$ -module $\Lambda_{\mathcal{L}}$. There exists a unique (up to a scalar) horizontal $u^{[i]} \in \Lambda_{\mathcal{L}^{[i]}}$ of the form

$$u^{[i]} = \sum_{r_1 \leq \dots \leq r_i} v_{r_1 \dots r_i}^{[i]} \eta^{(r_1)} \wedge \dots \wedge \eta^{(r_i)}$$

with $v_{n-i, \dots, n-1}^{[i]} \neq 0$. (Notice that $\eta^{(n-i)} \wedge \dots \wedge \eta^{(n-1)} \neq 0$ in $\Lambda_{\mathcal{L}^{[i]}}$.) With the induced pairing,

$$\mathcal{L}^{[i]} \langle u^{[i]}, \eta \wedge \dots \wedge \eta^{(i-1)} \rangle = \langle u^{[i]}, \mathcal{L}^{[i]} \left(\eta \wedge \dots \wedge \eta^{(i-1)} \right) \rangle = 0.$$

Thus we have, after modifying by a scalar, $v_{n-i, \dots, n-1}^{[i]} = \beta^i wr_i \in K[[t]]^\times$. Therefore, by subtracting an element in U_{i-1} , we obtain u_i satisfying (11) and (12). \square

Corollary 1.5 *Suppose \mathcal{L} in (1) is Calabi-Yau. With notations in Thm 1.4, we have $\langle U_{i-1}, U_{n-1-i} \rangle = 0$ and $\tau_i = \tau_{n-i}$ for all $1 \leq i \leq n$.*

Proof. By the last assertion of Thm 1.4, we see that $K(t) \otimes_{K[t]} U_i$ are the only possible $K((t))[\theta]$ -submodules of $K(t) \otimes_{K[t]} \Lambda_{\mathcal{L}}$. Since U_i is stable under θ , so is its orthogonal complement U_i^\perp . Thus by a rank counting, $U_i^\perp = U_{n-2-i}$.

Since u_i lies in Fil^i and $\langle \text{Fil}^i, \text{Fil}^{n-i} \rangle = 0$, we have $\langle u_i, u_{n-1-j} \rangle = 0$ if $i \neq j$ and $\langle u_i, u_{n-1-i} \rangle$ are non-zero constants. Consequently,

$$\begin{aligned} 0 &= \langle u_i, u_{n-i} \rangle' \\ &= \tau_i \cdot \langle u_{i-1}, u_{n-i} \rangle + \tau_{n-i} \cdot \langle u_i, u_{n-1-i} \rangle. \end{aligned}$$

Since $\tau_i(0) = 1$ for all i by definition, the second assertion follows. \square

2 Degenerations

For the definitions and basic properties of logarithmic structures, see [3].

Fix a base field k . Consider a flat projective pencil $\pi : X \rightarrow \mathbb{P}^1$ whose generic fiber is smooth. We further assume that each singular fiber of π is a union of reduced divisors with normal crossings. For such a pencil π , we equip X and \mathbb{P}^1 with the natural logarithmic structures associated to the union of the singular fibers (which is a reduced normal crossing divisor on X) and the critical values (which form a reduced divisor on \mathbb{P}^1), respectively. Then π is log-smooth. Denote by $\omega^i = \omega_{X/\mathbb{P}^1}^i$ the sheaf on X of relative differential i -forms with log poles with respect to the log structures.

Now suppose that the generic fiber of π is an absolutely irreducible Calabi-Yau variety of dimension $m \geq 1$. We will call such a π a *nice pencil of Calabi-Yau varieties of dimension m* . Then the sheaf $\pi_* \omega^m$ is an invertible sheaf on \mathbb{P}^1 . Suppose there exists a locally direct factor \mathcal{M} of $R^m \pi_* \omega^\bullet$ of rank $(m+1)$ which contains $\pi_* \omega^m$ and is stable under the Gauss-Manin connection ∇ . Now suppose $k = \mathbb{C}$. Let $a \in \mathbb{P}^1(\mathbb{C})$ be a \mathbb{C} -valued point and let N denote (the logarithm of) the local monodromy around a . Then N acts on the stalk \mathcal{M}_a at a and is nilpotent. Notice that if $N^m \neq 0$ on \mathcal{M}_a , then \mathcal{M} must be the unique irreducible locally direct factor of $R^m \pi_* \omega^\bullet$ which contains $\pi_* \omega^m$ and is stable under ∇ .

We make the following working definition, which is a special variant of being Hodge-Tate in the sense of Deligne ([2, §6]).

Definition. With notations and assumptions as above, we call \mathcal{M} *totally degenerate* at a if $N^m \neq 0$ on \mathcal{M}_a . We will abuse the notation by saying that the fiber at a of the family π is totally degenerate if such an \mathcal{M} exists. We call \mathcal{M} the *degenerate factor* of π in this case.

Lemma 2.1 *Let $\pi : X \rightarrow \mathbb{P}^1$ over \mathbb{C} be a nice pencil of Calabi-Yau varieties of dimension m as above with a totally degenerate fiber at 0. Then the Poincaré pairing is non-degenerate around 0 on the degenerate factor \mathcal{M} .*

Proof. Let η be a local basis of sections of $\pi_*\omega^m$ near 0. By assumption, we have, for $0 \leq i \leq m$,

$$\eta^{(i)} \in \text{Fil}^{m-i} \setminus \text{Fil}^{m-i+1}$$

and they form a local basis of \mathcal{M} at 0. Here $\eta^{(i)} = (\nabla(\theta))^i \eta (= N^i \eta$ at 0). Since $(\text{Fil}^i)^\perp = \text{Fil}^{m+1-i}$, the cup-product $\gamma := \langle \eta, \eta^{(m)} \rangle$ is an invertible function near 0. By Lemma 1.3, the assertion follows. \square

Corollary 2.2 *Let $\pi : X \rightarrow \mathbb{P}^1$ over $K \subset \mathbb{C}$ be a nice pencil of Calabi-Yau varieties of dimension m with a totally degenerate fiber at 0. Let η be a local basis of sections of $\pi_*\omega^m$ at 0 and let \mathcal{L} be the Picard-Fuchs operator of η . Then \mathcal{L} is a Calabi-Yau differential equation with respect to the parameter t of order $(m+1)$.*

Proof. The self-adjointness of \mathcal{L} follows from Lemma 2.1. The validity of condition (N) is obvious by the total degeneracy assumption. \square

3 Mod p and p -adic aspects

In this section, we fix a prime p and suppose $p > m$. Let K be a finite unramified extension of \mathbb{Q}_p with ring of integers W and residue field k . Let $\pi : X \rightarrow \mathbb{P}^1$ be a nice pencil of Calabi-Yau varieties of dimension m over K with a totally degenerate fiber at 0. Let ω^i be the sheaf on X of relative differential i -forms with log-poles along the logarithmic structure. We fix a basis η near 0 of sections of $\pi_*\omega^m \subset R^m\pi_*\omega^\bullet$ and let \mathcal{L} be the Picard-Fuchs operator of η as before. We call that π has *nice* reduction if π has a flat model over W such that (i) the reduction $\bar{\pi} : \bar{X} \rightarrow \mathbb{P}^1$ over k is also a nice pencil of Calabi-Yau varieties, and (ii) the two log structures of π and $\bar{\pi}$ are induced from a smooth log structure on the flat model.

(a) *The p -adic input*

Lemma 3.1 *Suppose the pencil π has nice reduction and the degenerate factor \mathcal{M}_0 at 0 is stable under (a lifting of) the absolute Frobenius. Then the Frobenius action on \mathcal{M}_0 is ordinary and consequently \mathcal{M} is generically ordinary.*

Proof. (Cf. [9, Thm 2.2].) Since \mathcal{M} is totally degenerate at 0, the Hodge structure on \mathcal{M}_0 is a successive extension of rank 1 Hodge structures of pairwise different weights. Since the Frobenius respects the Hodge and the weight filtrations (cf. [5, Remarques 3.28]), the result follows. \square

Lemma 3.2 *Suppose the pencil π over K with a totally degenerate fiber at 0 has nice reduction and the degenerate factor \mathcal{M}_0 at 0 is stable under Frobenius. Then there exists a non-zero constant $c \in W$ such that $c\beta F \in W[[t]]$.*

Proof. By Lemma 3.1 and [4, Prop 3.1.3], there is a non-zero element

$$u = \sum_{i=0}^m v_i \eta^{(i)}, \quad v_i \in W[[t]]$$

which is horizontal. By Thm 1.4, $v_m = c\beta F$ for some constant c . \square

Theorem 3.3 *Let π be a nice pencil of Calabi-Yau varieties of dimension m over \mathbb{Q} . Fix a basis near 0 of sections of $\pi_*\omega^m$ and let \mathcal{L} be the associated Picard-Fuchs operator. Let F be the formal solution of \mathcal{L} as in (7). Then $F(t) \in (1 + t\mathbb{Z}_p[[t]])$ for all p sufficiently large. In particular, $F(t) \in \mathbb{Z}[1/N][[t]]$ for some integer $N \neq 0$.*

Proof. (Cf. [6, Conj 2.1].) By the Lemma above and its proof, we see that except for a finite number of primes, $\beta F \in \mathbb{Z}_p[[t]]$ for all the remaining p such that the pencil π has nice reduction. By Lemma 1.3, $\beta(t)$ is rational over \mathbb{Q} and hence is in $(1 + t\mathbb{Z}_p[[t]])$ for almost all p . Thus the assertion follows. \square

(b) *The Hasse invariant and the differential equation*

Assume that π has nice reduction $\bar{\pi}$. Let $\bar{\omega}^i$ be the sheaf on \bar{X} of relative differential i -forms with log-poles. We choose η that can be extended to the flat model of π and assume the following condition is satisfied:

(R) Non-degeneracy of the reduction of η at the origin:

$$\langle \eta, \eta^{(m)} \rangle \in W^\times + tW[[t]].$$

Let σ be the lifting to $W[[t]]$ of the absolute Frobenius defined by sending t to t^p . For any $x \in W[[t]]$, denote by x^σ the image under σ .

Consider the adjoint morphism

$$V : \bar{\pi}_*\bar{\omega}^m \rightarrow \sigma^*\bar{\pi}_*\bar{\omega}^m$$

of the absolute Frobenius with respect to the cup-product on $R^m\pi_*\bar{\omega}^\bullet$. Notice that V is horizontal since the Frobenius and the cup-product are. Represent V by \mathfrak{H} defined by

$$V(\eta) = \mathfrak{H} \cdot \sigma^*\eta. \tag{13}$$

With our choice of the parameter, \mathfrak{H} is an element in $k(t)$. We call \mathfrak{H} the *Hasse invariant* of the family $\bar{\pi}$ (with respect to η). Notice that $\mathfrak{H}(0)$ is well-defined and non-zero at $t = 0$ by Lemma 3.1.

Proposition 3.4 *We have $\mathcal{L}\mathfrak{H} = 0$. That is, with respect to the parameter t , the Hasse invariant \mathfrak{H} is a rational solution of \mathcal{L} in characteristic p .*

Proof. By the very definition, $\nabla(\mathcal{L})\eta = 0$ and $\nabla\sigma^*\eta = 0$. Since V is horizontal, we see that $\mathcal{L}\mathfrak{H} = 0$ by applying $\nabla(\mathcal{L})$ to the equation (13). \square

If (R) is satisfied, then by Lemma 1.3, 3.2 and their proofs, $F \in W[[t]]$. Let

$$\mathcal{H}(t) = \frac{F(t)}{F(t)^\sigma} \tag{14}$$

regarded as a formal power series over W . Let $\bar{\mathcal{H}}(t)$ be the reduction mod p of $\mathcal{H}(t)$.

Proposition 3.5 *Suppose π satisfies the condition (R) above. Then we have the following.*

- (i) $\bar{\mathcal{H}}(t) \in (1 + tk[[t]]) \cap k(t)$.
- (ii) *The function $\bar{\mathcal{H}}$ satisfies $\mathcal{L}\bar{\mathcal{H}} = 0$.*

Proof. With the notations in Thm 1.4, the element $u_0 = \beta F \eta^{(m)} + \dots$ is a local generator of the submodule U_0 of horizontal sections of the degenerate factor \mathcal{M} (cf. the proof of Lemma 3.2). Then U_0 is a unit-root F -(iso)crystal ([4, Prop 3.1.3]) and hence by [4, 4.1.9], the function $\beta F / (\beta F)^\sigma$ is a lifting of an element in $k(t)$. Thus the first assertion follows.

Formally we have

$$F(t) = \frac{F(t)}{F(t)^\sigma} \frac{F(t)^\sigma}{F(t)^{\sigma^2}} \dots = \mathcal{H}(t) \mathcal{H}(t)^\sigma \dots$$

Applying the differential operator \mathcal{L} , we have

$$\begin{aligned} 0 = \mathcal{L}F(t) &= \mathcal{L}(\mathcal{H}(t) \mathcal{H}(t)^\sigma \dots) \\ &\equiv (\mathcal{L}\mathcal{H}(t)) (\mathcal{H}(t)^\sigma \dots) \\ &\equiv (\mathcal{L}\bar{\mathcal{H}}(t)) (\bar{\mathcal{H}}(t)^\sigma \dots) \pmod{p}. \end{aligned}$$

Thus the rational function $\bar{\mathcal{H}}(t)$ is a solution of \mathcal{L} in characteristic p . □

Proposition 3.6 *Assume condition (R) is fulfilled. Let $c = \mathfrak{H}(0)$. We have $\mathfrak{H} = c\bar{\mathcal{H}}$ regarded as rational functions of t .*

Proof. Over a non-empty open subset of \mathbb{P}^1 , both functions \mathfrak{H} and $c\bar{\mathcal{H}}$ represent the absolute Frobenius action on η ([4, 4.1.9]; cf. the proof of Prop 3.5(i)). Thus the assertion follows. □

Corollary 3.7 *Assume that condition (R) is fulfilled and $\pi_* \omega^m = \mathcal{O}(1)$. Suppose $\eta \in \Gamma(\mathbb{P}^1, \pi_* \bar{\omega}^m)$ and let $c = \mathfrak{H}(0)$. Then $c\bar{\mathcal{H}}(t) = \mathfrak{H}(t) \equiv cF^{<p}(t) \pmod{p}$, where $F^{<p}(t)$ is the truncation of $F(t)$ up to degree $(p-1)$.*

Proof. Under the assumptions, $\mathfrak{H}(t)$ is a polynomial of degree $< p$. □

Remark. The statement of the corollary is equivalent to the mod p case of the Dwork congruences of the coefficients $\chi(n)$ of $F(t) = \sum \chi(n)t^n$ in [6, §2.3]. Indeed let $\nu = \sum_{i \geq 0} \nu_i p^i$, $0 \leq \nu_i < p$, be the p -adic expansion of an integer $\nu \geq 0$. Then Cor 3.7 implies

$$\chi(\nu) \equiv \prod_{i \geq 0} \chi(\nu_i) \pmod{p},$$

which is equivalent to the mod p case.

On the other hand, if $\pi_* \omega^m = \mathcal{O}(n)$ for some $n > 1$, then $\bar{\mathcal{H}}$ is of higher degree. This may be used to detect the degree of $\pi_* \omega^m$ from the period F .

(c) *The higher congruences*

We keep the assumptions and notations in Lemma 3.1; assume that π has nice reduction $\bar{\pi}$ and (R) is fulfilled.

In this subsection, we are interested in understanding the higher Dwork congruences geometrically. For this purpose, we suppose that $\pi_*\omega^m = \mathcal{O}(1)$ over W and $\eta \in \pi_*\omega^m(\mathbb{P}^1)$ is globally defined. Thus β^{-1} , which equals $\langle \eta, \eta^{(m)} \rangle$ up to a multiplicative constant, is in $W[t]$.

We use the following notations. Let $\mathcal{H}_0 = F^{<p}(t)$. Denote by $\Delta_0 = \text{Spec } k[t]/\bar{\mathcal{H}}$ the *Hasse locus* of the pencil $\bar{\pi}$. Let $\Delta_\infty = (\mathbb{A}_W^1)_{/\Delta_0}$ be the completion of the affine line \mathbb{A}_W^1 along Δ_0 . Define

$$R_\infty = \varprojlim_n W[t, \mathcal{H}_0^{-1}]/(p^{n+1}),$$

and let

$$S_\infty = \text{Spf } R_\infty.$$

Under this circumstance, $\mathcal{H} \in R_\infty$ is a section of $\mathcal{O}(p-1)$ with $\mathcal{H} \equiv \mathcal{H}_0 \pmod{p}$ (cf. Cor 3.7 and [4, 4.1.9]; locally \mathcal{H} defined in (14) represents the absolute Frobenius on $\beta^{-1}u$ up to a multiplicative constant, where u occurs in the proof of Lemma 3.2). On the other hand, the unit-root part of \mathcal{M} defines a (multiplicative) formal group over S_∞ .

We make a further assumption:

- (E) The above formal group over S_∞ can be extended to a formal group G over the completion $(\mathbb{P}_W^1)_{/\mathbb{P}_k^1}$.

For example ([9]), the Dwork pencil has certain degenerate fibers; however the attached formal group can be extended to the whole $(\mathbb{P}_W^1)_{/\mathbb{P}_k^1}$ in an explicit way by writing down its logarithm (see [9, Prop 5.2]).

Now suppose (E) holds. Represent G by a formal group law over the affine part $(\mathbb{A}_W^1)_{/\mathbb{A}_k^1}$ with logarithm

$$l(\tau) = \sum_{n=1}^{\infty} \alpha(n) \frac{\tau^n}{n}, \quad \alpha(n) \in \varprojlim W[t]/(p^{i+1}),$$

which is normalized by $\alpha(p^n) \equiv \mathcal{H}\alpha(p^{n-1})^\sigma \pmod{p^n}$ for all $n \geq 1$. Write $F(t) = \sum \chi(n)t^n$. For any $1 \leq m < p$ and $s \geq 0$, consider the truncation of $F(t)$

$$F_{(mp^s)} = \sum_{n=(m-1)p^s}^{mp^s-1} \chi(n)t^n.$$

Lemma 3.8 *With notations as above, let $\{a_n\}_{n=0}^\infty, a_n \in W[t]$, be a sequence with $a_0 \in W^\times$ and $a_{n+1} \equiv \mathcal{H}a_n^\sigma \pmod{p^{n+1}}$ for all $n \geq 0$. Then we have the following.*

- (i) *For any $b \in W[t]$ with $b \equiv a_1 \pmod{p}$, there exists a sequence $\{b_n\}_{n=1}^\infty, b_n \in W[t]$, such that $b_1 = b$ and $b_{n+1} \equiv \mathcal{H}b_n^\sigma \pmod{p^{n+1}}$ for all $n \geq 1$.*

(ii) Fix a positive integer n . For any $c \in W[t]$ with $c \equiv a_n \pmod{p^n}$, there exists an element $\tilde{c} \in W[t]$, unique modulo p^{n+1} , such that

$$a_{n+1} \cdot c^\sigma \equiv a_n^\sigma \cdot \tilde{c} \pmod{p^{n+1}}.$$

Proof. (i) Indeed, if $b = a_1 + p\delta$, one checks that the sequence of elements

$$b_n = a_n + p \cdot a_{n-1} \delta^{\sigma^{n-1}}, \quad n \geq 2$$

(with $b_1 = b$) satisfies the requirement.

(ii) The assumption implies $a_{n+1} \equiv a_1^{1+p+\dots+p^n} \pmod{p}$. Write $c = a_n + p^n \alpha$. We are asked to solve ε in

$$a_{n+1} (a_n + p^n \alpha)^\sigma \equiv a_n^\sigma (a_{n+1} + p^n \varepsilon) \pmod{p^{n+1}},$$

which is equivalent to solve

$$a_n^\sigma \varepsilon \equiv a_{n+1} \alpha^\sigma \pmod{p}.$$

The latter has a unique solution $a_1 \cdot \alpha^\sigma$ modulo p . □

Theorem 3.9 *With notations as above and under the assumption (E), the formal power series*

$$\tilde{l}(\tau) = \sum_{\substack{s \geq 0 \\ 1 \leq m < p}} F_{(mp^s)} \frac{\tau^{mp^s}}{mp^s}$$

is the logarithm of a formal group law \tilde{G} over $(\mathbb{A}_W^1)_{/\mathbb{A}_k^1}$. Moreover, \tilde{G} is strictly isomorphic to G defined above over S_∞ .

Proof. We shall show that the sequence $\{F_{(mp^s)}\}$ satisfies the congruence property

$$F_{(mp^s)} \equiv \mathcal{H} \cdot F_{(mp^{s-1})}^\sigma \pmod{p^s}. \quad (15)$$

Consequently the theorem follows by [7, Thm (A.8), (A.9)].

For $s = 1$, the statement is a consequence of Cor 3.7.

Away from Δ_∞ , we have

$$\alpha(m, s) := \frac{\alpha(mp^s)}{\alpha(mp^{s-1})^\sigma} \in R_\infty$$

and $\alpha(m, s) \equiv \mathcal{H} \pmod{p^s}$ for all $s \geq 1$. To shorten the notation, first assume that $\mathcal{H} \pmod{p}$ is exactly of degree $(p-1)$. Now since $\alpha(m, s)$ represents the absolute Frobenius modulo p^s , which is a section of the sheaf $\mathcal{O}(p-1)$ of \mathbb{P}^1 over the affine open $\text{Spec } W[t, \mathcal{H}_0^{-1}]/(p^s)$, we have

$$\alpha(m, s) \equiv \frac{A}{\mathcal{H}_0^n} \pmod{p^s} \quad (16)$$

for some n and $A \in W[t]/(p^s)$ of degree $(n+1)(p-1)$.

On the other hand, applying Lemma 3.8(i) with $a_n = \alpha_{1,n}$ and $b = F_{(mp)}/F_{(m)} (\in W[t]$ with invertible constant term), we obtain a sequence $\{b_n\}$ such that, for $s \geq 1$,

$$\alpha(m, s) \equiv \frac{b_s}{b_{s-1}^\sigma} \equiv \mathcal{H} \pmod{p^s}. \quad (17)$$

(Here we set $b_0 = F_{(m)}$.) By the first congruence in (17) together with (16), we see that $\deg b_s = mp^s - 1$. Now by a simple induction argument, the second congruence in (17) together with Lemma 3.8(ii) (for $\{b_n\}$) imply that one can take $b_s = F_{(mp^s)}$ for all $s \geq 0$.

In general, we represent the occurred finite degree elements as homogeneous elements in $W[t_0, t_1]$ (e.g., $t_0^{mp^s-1} \alpha(t_1/t_0)$ instead of α) to obtain that $\deg b_s < mp^s$ and then repeat the argument above. This completes the proof. \square

Corollary 3.10 *Write $F(t) = \sum \chi(n)t^n$. For all non-negative integers ν, m, s with $0 \leq \nu < p$, we have*

$$\chi(\nu + mp^{s+1}) \equiv \chi(\nu)\chi(mp^s) \pmod{p^{s+1}}.$$

Proof. One checks readily that it suffices to establish the congruences for $1 \leq m < p$. Assuming m is in this range. The theorem above (see (15)) implies that, for all $s \geq 0$,

$$F \cdot F_{(mp^s)}^\sigma \equiv F_{(mp^{s+1})} \cdot F^\sigma \pmod{p^{s+1}}. \quad (\kappa_{m,s})$$

For $m = 1$, by comparing the coefficients of $t^{\nu+p^{s+1}}$ in both sides of the equation $(\kappa_{1,s})$, we get

$$\sum_{j=0}^{p^s-1} \chi(\nu + p^{s+1} - pj)\chi(j) \equiv \sum_{j=0}^{p^s-1} \chi(\nu + pj)\chi(p^s - j) \pmod{p^{s+1}}.$$

By canceling common terms on the two sides of the congruence, the assertion (corresponding to $j = 0$) follows immediately. For $m > 1$, by comparing the coefficients of $t^{\nu+mp^{s+1}}$ in both sides of equations $(\kappa_{m,s})$ and $(\kappa_{m-1,s})$, one obtains the desired congruences by induction on m . \square

Question. Is there a geometric/homological interpretation of the general Dwork congruences?

A Hadamard products

Here we describe how to obtain the unit roots precisely for certain Hadamard products considered in [6, §3], . The only missing piece in loc.cit. is to determine the constant ϵ_4 in Prop 2.7 there. To do this, we study the Frobenius action on the cohomology of the totally degenerate fiber by applying the weight spectral sequence in [5]. For references of Hadamard products and examples of pencils of elliptic curves we discuss here, see [6, §§3.1 and 3.2].

Let $X, Y \rightarrow \mathbb{P}^1$ be two pencils of elliptic curves over a finite field k of characteristic p with totally degenerate fibers X_0, Y_0 at 0, respectively. We assume that X_0, Y_0 are

strictly normal crossing divisors. Let ξ_1 and ξ_2 be local bases at 0 of horizontal sections of the relative H_{cris}^1 of X and Y over \mathbb{P}^1 , respectively. Then ξ_i are eigenvectors of the relative Frobenius. Denote by c_i the corresponding eigenvalues. Then $c_1 = 1$ if the degenerate curve X_0 is of split multiplicative type over k ; $c_1 = -1$ if X_0 is non-split. The corresponding statement for c_2 is similar.

Geometrically, the Hadamard product comes from the following commutative diagram with squares 1 and 2 being Cartesian:

$$\begin{array}{ccccc}
 Z & \longrightarrow & \widetilde{X \times Y} & \longrightarrow & X \times Y \\
 \downarrow & & \downarrow & \textcircled{2} & \downarrow \\
 & & \textcircled{1} \quad \mathbb{P}^1 \times \mathbb{P}^1 & \xrightarrow{b} & \mathbb{P}^1 \times \mathbb{P}^1 \\
 & & \downarrow & & \downarrow \pi \\
 0 & \longrightarrow & \mathbb{P}^1 & \xlongequal{\quad} & \mathbb{P}^1.
 \end{array}$$

Here π is the (coordinate-wise) multiplication; b is the blow-up of $\mathbb{P}^1 \times \mathbb{P}^1$ along $(\infty, 0)$ and $(0, \infty)$.

Over \bar{k} , let $Z^{(i)}$ be the disjoint union of all possible intersections of i distinct irreducible components of Z . Then we have the weight spectral sequence ([5, 3.23]):

$$E_1^{-j, i+j} = \bigoplus_{r \geq 0, r \geq -j} H_{cris}^{i-j-2r}(Z^{(1+j+2r)}/W)(-j-r) \implies H^i(Z^\times/W^\times), \quad (18)$$

which degenerates at E_2 modulo torsion ([5, Th 3.32]). Here W is the ring of Witt vectors of k and the target $H^i(Z^\times/W^\times)$ is the i -th logarithmic crystalline cohomology of Z .

Now assume that X and Y are ordinary and have trivial crystalline cohomology groups of odd degrees. Then the weights of the E_1 -terms are all integers and the non-zero terms of E_1 appear only when the weights are even.

Lemma A.1 *With assumptions as above, let ξ be the Hadamard product of ξ_i and c the eigenvalue of the relative Frobenius action on ξ . Then $c = c_1 c_2$.*

Proof. Let $d : E_1^{2,0} \rightarrow E_1^{3,0}$ be the boundary map in (18). Then c represents the relative Frobenius action on the cokernel of d . Let K be the field of fractions of W . By [5, Lemme 5.2], we see that $(\text{coker } d) \otimes K$ is 1-dimensional and the Frobenius acts on ξ as the product of its actions on ξ_i . Thus the statement follows. \square

References

- [1] G. Almkvist and W. Zudilin, Differential equations, mirror maps and zeta values. *Mirror symmetry V*, 481-515, AMS/IP Stud. Adv. Math., 38, Amer. Math. Soc., Providence, RI, 2006.
- [2] P. Deligne, Local behavior of Hodge structures at infinity. *Mirror symmetry II*, 683-699, AMS/IP Stud. Adv. Math., 1, Amer. Math. Soc., Providence, RI, 1997.

- [3] K. Kato, Logarithmic structures of Fontaine-Illusie. *Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988)*, pp. 191-224, Johns Hopkins Univ. Press, Baltimore, MD, 1989.
- [4] N. M. Katz, Travaux de Dwork. *Séminaire Bourbaki, 24ème année (1971/1972)*, Exp. No. 409, pp. 167-200. Lecture Notes in Math., Vol. 317, Springer, Berlin, 1973.
- [5] A. Mokrane, La suite spectrale des poids en cohomologie de Hyodo-Kato. *Duke Math. J.* 72 (1993), no. 2, 301-337.
- [6] K. Samol and D. van Straten, Frobenius polynomials for Calabi-Yau equations. *Commun. Number Theory Phys.* 2 (2008), no. 3, 537-561.
- [7] J. Stienstra and F. Beukers, On the Picard-Fuchs equation and the formal Brauer group of certain elliptic K3-surfaces. *Math. Ann.* 271 (1985), no. 2, 269-304.
- [8] M. van der Put and M. F. Singer, *Galois theory of linear differential equations*. Grundlehren der Mathematischen Wissenschaften, 328. Springer-Verlag, Berlin, 2003.
- [9] J.-D. Yu, Variation of the unit root along the Dwork family of Calabi-Yau varieties. *Math. Ann.* 343 (2009), no. 1, 53-78.